

*Reports from the Environmental Archaeology Unit, York 2000/04, 51 pp.*

**Assessment of biological remains from 41-49 Walmgate York  
(site code 1999.941)**

by

Cluny Johnstone, John Carrott, Allan Hall, Harry Kenward and Darren Worthy

**Summary**

*Excavations carried out at 41-49 Walmgate, York, as part of the Time Team Live programme, yielded a total of 34 samples and a single box of bone and shell. Of the samples, 22 (SRS and BS samples) were processed on site and eight GBA samples were processed at the EAU. Analysis was limited to Anglo-Scandinavian and medieval deposits with secure dating evidence.*

*Preservation of biological remains within these deposits was exceptional in a number of ways, particularly the presence of dried (and not rewetted) plant remains and charred insects in association with waterlogged material. The range of plant and insect species was very similar to that observed in Anglo-Scandinavian deposits in other areas of York. The species present were indicative of the interiors of buildings, including moist floor environments and possibly also thatch. Dyeplants and invertebrates associated with wool were present in several deposits, as in many Anglo-Scandinavian deposits from York, suggesting textile processing in the vicinity.*

*The shell and vertebrate assemblages, whilst being well preserved, were too small to provide any useful insights into the economy of the site, other than that they mainly represent food waste.*

*The exceptional preservation of all bioarchaeological remains from this site and the quantity of information gained from such a small scale intervention, has once again highlighted the potential of deposits in this area for yielding a wealth of information on the environment and economy of Anglo-Scandinavian York. It would be expected that a large scale excavation in this area would generate bioarchaeological data on a scale similar to the work at 16-22 Coppergate.*

**KEYWORDS:** 41-9 WALMGATE; YORK; ASSESSMENT; PLANT REMAINS; CHARRED REMAINS; MICROFOSSILS; PHYTOLITHS; DIATOMS; INTESTINAL PARASITE EGGS; *TRICHURIS*; INSECTS; SHELLFISH; OYSTERS; VERTEBRATE REMAINS; FOOD; DYEPLANTS

Authors' address:

Palaeoecology Research Services  
Environmental Archaeology Unit  
Dept. of Biology  
University of York  
PO Box 373  
York YO10 5YW

Prepared for:

York Archaeological Trust  
Cromwell House  
13 Ogleforth  
York  
YO1 7FG

Telephone: (01904) 433846/434475/434487  
Fax: (01904) 433850

3rd March 2000

# Assessment of biological remains from 41-49 Walmgate, York (site code 1999.941)

## Introduction

Excavations at 41-49 Walmgate (NGR SE607515) took place over the weekend of 3th-5th September 1999, as part of the 'Time Team Live' television programme for Channel 4. The excavation was undertaken by York Archaeological Trust. An on-site sieving programme (using both coarse-riddling and bulk-sieving) was undertaken by staff from the EAU and the Archaeological Resource Centre, York. Samples were also taken for further analysis at the EAU.

The Time Team excavations included the re-excavation of a trench dug in 1990, by York Archaeological Trust, which revealed well-preserved deposits just under the ground surface. Work carried out on deposits recovered from that excavation (Carrott *et al* 1991) showed the potential for the recovery of useful quantities of well preserved biological remains.

Samples and vertebrate material from the 1999 intervention were recovered from layers dating to the Anglo-Scandinavian, medieval and post-medieval periods. Six phases of activity were identified and are dated as follows:

- Phase 1 - Anglo-Scandinavian
- Phase 2 - 11th-12th Century
- Phase 3 - ?12th-14th Century
- Phase 4 - Late medieval
- Phase 5 - Late medieval / post-medieval
- Phase 6 - Modern

## Methods

### *Sediment samples*

Descriptions of the lithologies of all of the samples were recorded using a standard *pro forma*. Site-riddling (coarse-sieving) was undertaken on site using a riddling frame with a 10 mm mesh, after disaggregation in a cement-mixer. Bulk-sieving was also performed on site using a modified S raf tank with a 1mm mesh and 1mm washover sieves. In addition, 'GBA' samples (*sensu* Dobney *et al.* 1992) were taken; these were processed at the EAU, following procedures of Kenward *et al.* (1980; 1986), for the recovery of plant and insect remains.

The samples were examined for the eggs of parasitic nematodes using the 'squash' technique of Dainton (1992).

### **Plant remains and other components of the GBA and BS samples**

Plant remains were examined in the dried bulk-sieving (BS) residues and washovers, and wet GBA residues and flots. A semi-quantitative record of the abundance of taxa was recorded using a three-point scale for the BS material (and for some 'finds' from BS residues resulting from on-site sorting), and a four-point scale for the GBA samples. At the same time, records were made of all other components of

these fractions—items such as bark, charcoal, wood, bone of various kinds, shell, and a variety of materials from occupation, such as pottery and brick/tile. The maximum size of most of these components was also noted. Records were made directly to a computer database form.

### **Insect remains**

Insects in paraffin flots from GBA samples were identified by comparison with modern reference material and using the standard works. Adult beetles and bugs, other than aphids and scale insects, were recorded fully quantitatively in order to put data on record for a synthesis project currently underway. A minimum number of individuals was estimated on the basis of the fragments present. Other invertebrate macrofossils were recorded semi-quantitatively using the scale described by Kenward *et al.* (1986) and Kenward (1992), using estimates for extremely abundant taxa.

Recording of the macro-invertebrates was essentially 'detailed' in the sense of Kenward (1992): many of the identifications were pushed further than is normal under 'scan' recording. Recording of the state of preservation of invertebrates followed Kenward and Large (1998), making use of the sheet illustrated in their fig. 2.

Data pertaining to invertebrate remains were transferred from a paper record to computer databases (using *Paradox* software) for analysis and long-term storage.

### **Shell**

A small quantity of shell recovered by both hand-collection and on-site riddling of sediment samples was submitted: a volume of approximately two litres, representing material from eight contexts. Five of these were poorly dated or modern and shell recovered from them was not recorded. Brief notes were made on the preservational condition of the shell and the remains identified to species where possible.

For oyster (*Ostrea edulis* L.) shell, additional notes were made regarding: numbers of left and right valves; evidence of the oysters having been opened using a knife or similar implement; measurability of the valves; damage from other marine biota (polychaete worms and dog whelks); encrustation by barnacles.

### **Vertebrate remains**

For the hand-collected vertebrate remains, data were recorded electronically directly into a series of tables using a purpose-built input system and *Paradox* software. Subjective records were made of the state of preservation, colour of the fragments, and the appearance of broken surfaces ('angularity'). Additionally, semi-quantitative information was recorded for each context concerning fragment size, dog gnawing, burning, butchery and fresh breaks.

Where possible, fragments were identified to species or species group, using the reference collection at the Environmental Archaeology Unit, University of York. Fragments not identifiable to species were described as the 'unidentified' fraction. Within this fraction fragments were grouped into a number of categories: large mammal (assumed to be cattle, horse or large cervid), medium-sized mammal (assumed to be caprovid, pig or small cervid), bird, fish, small mammal and totally unidentifiable. As well as counts of fragments, total weights were recorded for all identifiable and unidentifiable categories.

Vertebrate remains from the samples were recorded in a similar manner.

## Results

The results of these investigations are presented in the following account, supported by Table 3 for plants, Tables 4-9 for invertebrates, Table 10 for shell and Tables 11-14 for vertebrates.

### *Sediment samples*

A list of samples, together with notes on their treatment, is given in Table 1. The material is discussed in phase order, earliest to latest.

#### Phase 1

**Context 1003** [?floor deposit]

**Sample 11/GBA** (2kg subsample and microfossil 'squash')

Moist, mid to dark grey-brown, unconsolidated (working soft), clay sandy silt. Very small stones (2-6mm) were present together with charcoal.

The microfossil 'squash' was mostly organic detritus with a little inorganic material and very many phytoliths. No eggs of intestinal parasitic nematodes were seen.

The residue and flots from this subsample were characterised by the presence of rather large amounts of charred herbaceous material including stems (mostly unidentified, but with some grass and/or cereal stem nodes and traces of trigonous stem fragments, perhaps bulrush or sea clubrush, *Scirpus lacustris/maritimus*) and some extremely well preserved rachis (ear-stalk) fragments of rye (*Secale cereale*). The latter, which were sometimes on lightly charred ('toasted') bore the characteristic rows of stiff hairs along their margins. Also present were fragments of the lemma (part of the flower) of rye with their distinctive coarse marginal cilia. Some of the charred plant remains exhibited a 'lumpy', irregular charred surface perhaps result of smoke-blackening rather than charring in fire, suggesting they may be from rye straw in thatch which was subsequently partly burnt in a fire. The presence of rather frequent charred seeds of comcockle *Agrostemma githago* is consistent with this interpretation.

For the rest, the residue included large amounts of charcoal and some uncharred wood chips, sand and daub, the last consistent with the presence of material from a building. There were also some small concretions of ashy material, containing fine charred herbaceous stems, similar to material found in quantity at the Middle-Late Saxon site at Flixborough, N. Lincolnshire (Hall, in preparation) though the precise mechanism whereby these concretions form is as yet unknown. Other seeds included a range of weeds (mostly charred) with traces of uncharred flax (*Linum usitatissimum*) capsule, and charred grains or spikelets of oats (*Avena sativa*), barley (*Hordeum*), rye and bread/club wheat (*Triticum aestivo-compactum*). The presence of traces of charred *Triglochin maritima* (sea arrow-grass) fruits also resonates with the evidence from Flixborough, where remains of plants from the salt-marsh habitat (though not *Triglochin* itself) were quite often encountered in ash-rich deposits.

Although a 2kg subsample of the deposit was processed, rather few invertebrates were recovered: 52 individuals of 32 beetle and bug taxa, modest numbers of mites, and smaller numbers of a few other groups. The main statistics of the beetle and bug assemblages (Table 4) were broadly similar to those for deposits at comparable sites, e.g. 16-22 Coppergate (Kenward and Hall 1995). The upper ranks of abundance included several taxa regarded traditionally as 'house fauna' (Kenward and Hall 1995, 662-7) and placed in a group typical of floors by statistical analysis (Group A, Carrott and Kenward 2000): *Xylodromus concinnus* (7 individuals); *Atomaria* sp. (4); *Cryptophagus* sp. (3); and *Crataraea suturalis* and *Lathridius minutus* group (2 each), with single individuals of various others falling in this category. Of the beetles and bugs, 44% (23) fell in Carrott and Kenward's core Group A (Table 8), and there were also two human fleas, so that there is little doubt that this was either a floor deposit or rich in floor

sweepings.

There may have been traces of foul matter present, as indicated by two *Platystethus arenarius*, but this very mobile species may have arrived accidentally with the various ‘outdoor’ elements recorded. On balance, a fairly clean floor is indicated.

Charred insect remains were—exceptionally—fairly abundant in this deposit. Those which could be recognised are listed in Table 9 (there were other burned remains which may have been badly damaged invertebrates). The remains seem to have been charred in the floor *in situ*, although they may just possibly have been reduced in burning roofing material.

**Context 1004** [?floor deposit]

**Sample 12/GBA** (1kg subsample and microfossil ‘squash’)

Moist, dark brown-black, stiff and compressed (working crumbly), slightly humic, very slightly sandy, clay silt. Very small stones (2-6mm) were present. Other inclusions were charcoal, wood and large mammal bone fragments.

The microfossil ‘squash’ was approximately half-and-half organic detritus and inorganic material with a few fungal spores and hyphae. No eggs of intestinal parasitic nematodes were seen.

There were moderate amounts of well preserved waterlogged (and sometimes also charred) plant remains in the residue and flots from this sample, the matrix consisting largely of small wood chips, bark, charcoal and herbaceous detritus. There were some uncharred oats (*Avena*) spikelets which looked as if they had dried and been rewetted (almost as if they were rather decayed modern specimens which had not long been buried). Some fragments of hazel (*Corylus avellana*) nutshell showed evidence of knife marks at the apex (cf. Kenward and Hall, 1995, fig. 19 1g, h). There were also traces of three plants likely to have been used in textile dyeing: stems fragments of the clubmoss *Diphysium complanatum*, a root fragment of madder (*Rubia tinctorum*) and a pod fragment of woad (*Isatis tinctoria*), all recorded regularly from Anglo-Scandinavian deposits in York (e.g. Kenward and Hall 1995).

Substantial numbers of beetles were recovered from this sample: 167 individuals of 77 taxa. Other invertebrates were moderately abundant. In addition to those mentioned below there were ‘many’ proctotrupoid wasps and mites, ‘several’ earthworm egg capsules, fly puparia and beetle larvae, and three frog hopper nymphs.

The beetles were somewhat mixed ecologically, but ‘house fauna’ taxa were numerically predominant: the most abundant species was *Xylodromus concinnus*, with 29 individuals, and there were ten *Lathridius minutus* group, eight *Crataraea suturalis*, six *Cryptophagus scutellatus*, and three each of *Ptinus ?fur* and *Atomaria nigripennis*. Carrott and Kenward’s (2000) core Group A accounted for 45% of the adult beetles (Table 8), so the interpretation of this deposit as a floor seems reasonable. There were also eight human fleas (*Pulex irritans*), which would have bred in floor litter. Many of the other species recorded would have found habitats in litter which was somewhat damp, but not objectionable to humans. There were, however, some beetles typical of rather more foul conditions, notably *Platystethus arenarius* (4), and *Anotylus nitidulus* and *Gyrophynus angustatus* (3 each). These may have been strays, or indicative of local (and perhaps very small) patches of foul matter.

The cleaning of wool or fleeces was indicated by two adult and one puparial sheep ked, *Melophagus ovinus*. The dung deposited in cleanings (‘dags’) may have attracted the ‘foul’ beetles mentioned above, and also the dung beetle *Aphodius ?prodromus*, of which there were two.

This deposit may have formed in a well-protected situation within a building, for the proportion of

'outdoor' insects (PNOB, Table 4) was low by comparison with samples from 16-22 Coppergate and other sites. This is supported to some extent by a provisional record of a head capsule of a newly-emerged individual of the very cryptic and synanthropic ground beetle *Sphodrus leucophthalmus*.

A single eel vertebra and vole/mouse (microtine/murine) astragalus were the only vertebrate remains recovered from this sample.

#### **Sample 13/BS**

See Sample 12 for description.

The small washover of about 100cm<sup>3</sup> from this sample, consisted mostly of woody debris but included some cereal chaff with grades between charred and uncharred, some of it rye rachis. It was noted during recording that this material might have been 'dry-preserved' (as in thatch) before being buried. The presence of saw-sedge (*Cladium mariscus*) and sedge (*Carex* spp.) nutlets with their perigynia (outer 'coats') perhaps adds weight to the interpretation that whole dried plants were becoming incorporated then buried into the deposit. As in Context 1003, there were fragments of rye lemma and rachis (and also some tentatively identified waterlogged rachis fragments), though these were not seen in the GBA subsample.

Vertebrate remains comprised two herring vertebra.

#### **Sample 14/BS**

See Sample 12 for description

A spot find of hazelnutshell was recovered from this sample. In addition, 'unidentified' vertebrate remains (weighing 9.5g) were recovered, falling into the large- and medium-sized mammal, bird, and completely unidentified categories. Preservation of the vertebrate remains appeared mixed.

#### **Context 1005** [black humic layer]

##### **Sample 6/SRS**

Moist, dark brown, soft, very slightly humic, very slightly sandy, clay silt. Stones up to 20mm were present, together with wood fragments.

A spot find of hazelnutshell was recovered from sample 6/SRS. Vertebrate remains comprised a single bird bone and three unidentified fragments (weighing 4.2g).

##### **Sample 8/BS**

See Sample 6 for description.

The washover from sample 8/BS yielded traces of clubmoss, woad, flax, and distinct wetland (fen/marsh) and cut grassland taxa. The frequent wood fragments were rather pale and spongy, giving an appearance of material that had been dried and rewetted at some point.

Although vertebrate remains were numerous (89 fragments, weighing 9.7 g), almost all (86) were very small and unidentifiable. The remaining three consisted of a single bird bone and two fish fragments. Overall preservation was good and colour dark brown. A few battered fragments were noted. More than half the fragments were burnt, which explains the high degree of fragmentation.

##### **Sample 10/GBA** (1kg subsample and microfossil 'squash')

See Sample 6 for description.

The microfossil 'squash' was mostly organic detritus with a little inorganic material and many phytoliths.

No eggs of intestinal parasitic nematodes were seen.

The large residue consisted of less than 10% by volume of mineral material (sand and grit), the rest being rather granular to flaky woody debris (including some chips) and bark fragments. The abundant small (<2mm) wood fragments may perhaps represent 'sawdust'. Together with fragments of clubmoss there was a single decayed wood pod fragment. Other plant remains included moderate numbers of well preserved seeds and fruits of a broad range weeds (especially stinking mayweed, *Anthemis cotula*) and fat hen (*Chenopodium album*), a variety of mosses, and traces of a few plants which might represent heathland/moorland habitats (brought, for example, in peat or turves?).

Insect remains were abundant and included 184 adult individuals of 88 beetle and bug taxa. Mites were very abundant, and there were moderate numbers of earthworm egg capsules and proctotrupoid wasps. The assemblage included two main components: 'house fauna', and species found in somewhat foul, but open-textured, matter such as stable manure (and probably dyeplant waste at 16-22 Coppergate). The first group included *Xylodromus concinnus* (20 individuals), *Aglenus brunneus* (14), *Cryptophagus scutellatus* (8), and three each of *Cryptophagus* sp., *Lathridius minutus* group, *Crataraea suturalis* and *Atomaria nigripennis*. There were also three human fleas. Beetles and bugs in Carrott and Kenward's core Group A contributed 42% of the assemblage, and an origin for this deposit within a building seems extremely likely.

The second ecological group was typified by *Ptenidium* sp. and *Leptacinus pusillus* (5 each), *Gyrohypnus fracticornis* (4), *Carpelimus fuliginosus* and *Oxytelus sculptus* (3 each), and some of the less common taxa. Species representing extremely foul conditions were rare, however, and it seems likely that most of the fauna exploited various aspects of the interior of a building. That this was fairly well closed is suggested by the relative rarity of 'outdoor' insects: single individuals of 17 taxa, many of which may in any case have been imported with water or cut wetland vegetation.

**Context 1014** [make up for floor]

**Sample 19/BS**

Just moist, light grey through mid grey and mid grey-brown to dark grey, stiff to crumbly (working soft), clay silt. Small (6-20mm) and medium-sized (20-60mm) stones and large mammal bone were present in this sample.

The very small washover (which included much undisaggregated sediment) contained flaky pale-coloured wood fragments with quite large numbers of well-preserved seeds, mainly weeds of various kinds. Flax, hop (*Humulus lupulus*) and clubmoss were all recorded. Fifty vertebrate fragments (weighing 60g) were recovered, including cattle, pig, bird, large mammal and unidentified remains. Preservation of the vertebrate remains was generally good, although a few battered fragments were noted. Colour was recorded as dark brown. Between a quarter and half the fragments were burnt.

**Sample 20/GBA** (1kg subsample and microfossil 'squash')

See Sample 19 for description

The microfossil 'squash' was mostly of inorganic material with some organic detritus and a few fungal hyphae and diatoms. No eggs of intestinal parasitic nematodes were seen.

This subsample yielded only a small residue consisting mainly of bark and wood fragments on the one hand, and concreted sediment (including 'root moulds'), grit and sand, on the other. There were low concentrations of moderately well-preserved seeds of which only stinking mayweed, fat hen and annual nettle (*Urtica urens*) were present in more than trace amounts. There were a few fragments of clubmoss and madder root, but otherwise most plant remains were from weeds or other wild-growing taxa.

The GBA subsample gave a modest-sized assemblage of adult beetles and bugs (44 taxa, 60 individuals) of rather mixed character. Of the other invertebrates noted, only mites and earthworm egg capsules were at all common. A quarter of the beetles (only 15 individuals) were from 'outdoor' forms, mostly probably representing open-air habitats on the site (ground beetles, dung beetles), a few probably further-travelled (e.g. the chafer *Hoplia philanthus*). There was a trace of house fauna (a fifth of the beetles, including three *Xylodromus concinnus*, two human fleas and a sheep ked), but no more than might be present in an outdoor deposit on an occupation site, brought to level a floor, and certainly no more than might be subsequently trampled into it.

### **Sample 21/SRS**

See Sample 19 for description.

This was an SRS sample processed on site; no biological remains were recovered.

### **Context 1034** [organic silt layer]

#### **Sample 31/GBA** (1kg subsample and microfossil 'squash')

Moist, mid-dark grey brown, crumbly, very humic silt with fine and coarse herbaceous detritus. Lumps of compressed, layered, sediment were present—possibly the same as the matrix but less disturbed. Wood and 'straw' were present. A modern contaminant in the form of a lollipop stick was noted!

The microfossil 'squash' was mostly of organic detritus with a little inorganic material and many phytoliths. No eggs of intestinal parasitic nematodes were seen.

This subsample gave a very large residue of about 450cm<sup>3</sup> of which all but a few tens of cm<sup>3</sup> were organic fragments, mainly wood with some bark and charcoal. The remainder was sand and gravel. There were also a few small (<5mm) lumps of sticky material with plant remains, including seeds, pressed into it which is thought to be sealing wax (it melted easily and with a characteristic smell and hardened to a dark brittle layer on a glass slide); this may also have been responsible for the rather pronounced yellowish colouration of the ethanol in the flot.

Seeds from this sample were mostly well preserved but not especially abundant (except for stinking mayweed, which was very frequent), though there were traces of root and root bark of madder and moderate numbers of stem fragments of clubmoss. *Cladium* nutlets with their perigynium present were also recorded, as were achenes of the sea aster, *Aster tripolium*, a plant characteristic of salt-marsh (and likely to have arrived, like the *Triglochin* from Samples 11, 13 and 20. There was also a distinct grassland component in Sample 31 which may have arrived in hay or stable manure.

The assemblage of adult beetles and bugs was rather limited in size (N = 65, S = 36), although a substantial range of other invertebrates was recorded. These included numerous mites, and 'several' fly puparia and proctotrupoid wasps (this group of wasps include many species parasitic in fly puparia, which are probably their main source in deposits such as these).

The beetles (and a few bugs) had an ecologically mixed character, and a quarter of the individuals were of 'outdoor' taxa. There were (limited) indications of 'house fauna' from four *Xylodromus concinnus* (the most abundant beetle), four human fleas, and a very few others, and hints of fouler matter from the oxyteline staphylinids (*Carpelimus pusillus* group and *Platystethus arenarius*, with three individuals each, being the most abundant).

Overall, this deposit appears to have formed in the open and to have received contributions from a wider range of sources. Subjectively, the insects offer a hint of the presence of material from heath or moor habitats, but the relevant taxa were too eroded and fragmentary for confident identification.



A notable record was a larval segment (bearing the characteristic spines illustrated by Peacock 1993) of the larder beetle *Dermestes lardarius*.

Three herring (*Clupea harengus*) vertebrae and two unidentified fish bones (weighing 0.2g) were recovered from this context.

**Context 1035** [fill of small cut 1036]

**Sample 32/GBA** (1kg subsample and microfossil 'squash')

Moist, light grey to mid to dark grey brown (and points in between), unconsolidated (working soft), slightly clay sandy silt. Fragments of rotted mortar and pot sherds were present in this sample.

The microfossil 'squash' was mostly of organic detritus with a little inorganic material and many phytoliths. A live soil nematode was also noted.

The moderate-sized to large residue of about 300cm<sup>3</sup> was less than one-third grit, sand and gravel, the rest being wood (including chips) and charcoal and bark fragments. Again, there were rather low concentrations of mostly well-preserved seeds, mainly of weeds but with remains of flax (seeds and capsule fragments), clubmoss and rye (rachis).

Mites were rather abundant, but other invertebrates, including beetles and bugs, were rare (N = 52, S = 36). The presence of nine *Aglenus brunneus*, four *Cryptophagus scutellatus*, three *Xylodromus concinnus*, and single individuals of some other taxa, suggests the presence of a house fauna component (and indeed half of the fauna fell into Carrott and Kenward's core Group A). There were a few remains of taxa suggesting foul conditions, but no evidence that any of them established breeding populations. There is nothing from the fauna to indicate whether this cut was located indoors, or was outdoors and received sediment containing house fauna.

Single gadid (cod family) and herring (*Clupea harengus*) vertebrae were recovered from this sample (total weight 1.2g).

**Context 1039** [organic silt layer]

**Sample 33/GBA** (1kg subsample and microfossil 'squash')

Moist, dark brown, layered and compressed to unconsolidated, silt with fine and coarse herbaceous detritus. Rotted wood, 'straw' and moss were present.

The microfossil 'squash' was mostly of organic detritus with a little inorganic material, some fungal spores and hyphae, and many phytoliths. No eggs of intestinal parasitic nematodes were seen.

There was a very large residue of about 750cm<sup>3</sup> of which only a few cm<sup>3</sup> were dense material (mainly bone and gravel), the rest woody detritus, mainly wood (including chips) and bark, with some charcoal. The >2mm fractions were particularly 'flaky' in appearance.

A wide range of mosses was present, all in small amounts, representing woodland habitats, especially bark and shaded rocks and trees. There were only a few seeds in the coarse fractions but these included a few well-preserved charred cereals (oats and barley) and moderate numbers of flax seeds; there were also traces of clubmoss and madder, and a single small fragment of dyer's greenweed (*Genista tinctoria*) stem. Particularly noticeable in the residue (and also in the flot) was the presence of fine herbaceous detritus with a 'grassy' or 'strawy' character, together with several grassland taxa represented by seeds or fruits. These may have originated in cut vegetation such as hay or straw, perhaps from stable manure. Traces of small concretions appeared to be of the kind seen in deposits rich in this kind of material rather than in deposits

of human faecal matter and, indeed, an additional test for parasite eggs on one fragment of concretion proved negative.

Beetles and bugs were not abundant (72 individuals of 50 taxa being recorded), but there were very large numbers of fly puparia, mostly Sphaeroceridae, and numerous mites. Although there was a small house fauna element (core Group A accounting for a fifth of the fauna), the more abundant taxa suggest formation under fairly foul conditions in the open. *Platystethus arenarius* was the most numerous species (with five individuals), followed by *Carpelimus pusillus* group and *Anotylus complanatus*—a group of species seen repeatedly in pits at 16-22 Coppergate. The rarer taxa continued this pattern, the presence of single individuals of four *Aphodius* dung beetles being notable. The family Sphaeroceridae includes many species whose immature stages are typically found under very foul conditions.

Three large mammal shaft fragments (weighing 17.8 g) were the only vertebrate remains recovered from this sample

### Phase 3

**Context 1031** [?dumping/build up in backyard]

#### **Sample 26/BS**

Moist, mid-dark brown, soft (working soft/plastic), very slightly sandy silty clay. Stones under 20mm and over 60mm were present. The only other inclusion noted was large mammal bone fragments.

The very small washover included traces of seeds of a few weeds, charred ?heather (*Calluna vulgaris*) root/twig fragments and flax seeds, and at least one charred 6-row barley (*Hordeum vulgare*) grain, as well as oats and bread/club wheat.

Of the 163 fragments of bone (weighing 66.4 g) recovered from this sample, 12 were identifiable. Species present included pig, mouse (Murine sp.), cod (*Gadus morhua*), eel (*Anguilla anguilla*) and herring (*Clupea harengus*). The unidentified fraction contained fragments of bird, fish, large- and medium-sized mammal bones together with 105 completely unidentifiable small pieces. Preservation was described as good and colour as dark brown. Angularity was recorded as variable with rounded, battered and spiky fragments all present.

#### **Sample 27/SRS**

See sample 26 for description.

Two small fragments (to 23mm) of ?oyster shell were recovered from this sample.

### Phase 4

**Context 1013** [fill of pit 1029]

#### **Sample 16/SRS**

Moist, mid-brown, crumbly (working sticky), slightly sandy silty clay. Stones ranging from 2-60+mm were present, together with brick/tile, mortar/plaster, pottery, marine mollusc shell and large mammal bone fragments.

A spot find of hazel nutshell was recovered from this sample.

The marine mollusc shell consisted of four left and two right oyster (*Ostrea edulis*) valves, fourteen fragments of oyster shell (largest to 40mm), and a single fragment of mussel (*Mytilus edulis*) shell. In

addition, three fragments of land snail (*Helix* sp.) were noted (probably all from one individual).

The vertebrate assemblage from this sample was quite extensive and reasonably well-preserved. A total of 278 fragments (weighing 949.8g) was recovered. Species present included cattle (8 fragments), pig (7) and caprovid (4), chicken (2), cod (*Gadus morhua*) (6 fragments), gadid (cod family) and haddock (*Melanogrammus aeglefinus*). The unidentified fraction included fragments of bird (7 fragments), fish (3), large (43) and medium-sized mammal (56) bones, in addition to 140 completely unidentified fragments.

**Context 1030** [fill of pit 1029]

**Sample 22/BS**

Moist, mid-dark grey-brown, crumbly to soft (working soft), very slightly sandy clay silt. Small stones (up to 20mm) were present together with brick/tile, wood and land snails.

The small wet residue of a few hundred cm<sup>3</sup> of woody debris was scanned very quickly: much of the material was rather spongy and floated, as if it had dried out and then been rewetted. Again, clubmoss was present, though it was very decayed. Two further taxa, recorded only from this sample amongst the assemblages from this site, were walnut (*Juglans regia*) and pea (*Pisum* cf. *sativum*).

A total of 342 bone fragments (weighing 77.3 g) was recovered from this sample. Species identified included pig, caprovid, gadid (cod family), eel (*Anguilla anguilla*) and herring (*Clupea harengus*). The unidentified fraction included fragments of bird, fish, large- and medium-sized mammal bones together with completely unidentified remains. Preservation was generally good, although a few battered fragments were noted and the colour was brown and dark brown. Burnt fragments formed 20-50% of the assemblage.

**Sample 23/SRS**

See Sample 22 for description.

Four small fragments (to 35mm) of oyster shell and two fragments (to 23mm) of mussel shell were recovered from this sample.

Two cod (*Gadus morhua*) vertebrae and a single haddock (*Melanogrammus aeglefinus*) vertebra (weighing together 5g) were recovered from this sample.

**Sample 24/GBA** (1kg subsample and microfossil 'squash')

See Sample 22 for description.

The microfossil 'squash' was approximately half-and-half inorganic material and organic detritus. A few fungal spores and phytoliths and a single *Trichuris* egg were noted. This last was somewhat pale but otherwise well-preserved (and measurable).

There was a moderate-sized residue of about 300cm<sup>3</sup> of which less than a third was mineral (sand, gravel, brick/tile); the rest mainly fine organic material (mainly very decayed wood). There was a little moss and some (?wool) yarn fragments but no suggestion of the presence of faecal material. There was also a possible peat/turf component. The presence of clubmoss in this late medieval context suggests that reworking of Anglo-Scandinavian material had taken place. Though preservation was generally good, the concentration of identifiable remains was low.

Only a small assemblage of adult beetles, and a single bug, was recovered (N = 36, S = 33) and other remains were rare apart from 'many' mites. Although no species was represented by more than two individuals, the beetle fauna had echoes of many recorded from pits at 16-22 Coppergate, the most

numerous species being *Carpelimus ?bilineatus*, *C. pusillus* group and *Platystethus arenarius* (only two of each, however). The presence of foul matter is supported by the proportion of taxa coded 'rf' (although the number of individuals is of course small, Table 5). Many of the remains were poorly preserved, and it is possible that some had completely decayed, but it seems equally likely that the deposit was buried too rapidly for a large insect population to develop following initial invasion by a range of species.

Although this deposit is dated to the late medieval period, it included no grain pests (a very constant signal in assemblages of such late date) and had a strongly 'Anglo-Scandinavian' character. This accords with the botanical evidence, strongly indicating that the layer consisted primarily of re-excavated earlier material.

Two herring (*Clupea harengus*) vertebrae, a single eel (*Anguilla anguilla*) vertebra and a bank vole (*Clethrionomys glareolus*) maxillary molar were recovered from this context. All were very well preserved.

## **Phase 6**

**Context 1000** [Backfill of previous excavation trench]

### **Samples 1, 2, 3 and 5**

Moist, mid to dark brown, crumbly to unconsolidated (working crumbly), very slightly clay, slightly silty sand. Stones ranging from 2-60+mm were common. Other inclusions noted were mortar/plaster, brick/tile, pottery, metal, glass, tarmac, plastic, charcoal and wood. Sample 5 contained humic material and fewer stones.

These were SRS and BS samples processed on site; no further action was taken, except that material from Sample 5/BS was checked briefly for the presence of dyeplants; traces of stem fragments of the clubmoss, *Diphasium complanatum* (recorded from most samples, see below), were recovered.

**Context 1001** [cleaning of previous trench]

### **Sample 4/SRS**

Moist, mid to dark slightly greyish brown, crumbly (working soft), slightly clay, sandy silt. Stones ranging from very small to large (2-60+mm) were common. Brick/tile, twigs and large mammal bone fragments were also present. Modern roots and rootlets, together with earthworm bioturbation were noted.

The sample was processed on site, no further action as the context was a cleaning layer.

**Context 1011** [machine excavated, mostly medieval finds]

### **Samples 7 and 17**

Moist, mid-brown, crumbly to soft (working soft), sandy clay silt. Stones ranging from 2 to 60+mm were present. Mortar/plaster and pottery were also present, whilst brick/tile fragments were common. Sample 17 was darker in colour and contained less rubble.

These were SRS and BS samples processed on site; no further action was taken as the context was machine-dug. A single spot find of hazel nutshell was noted from Sample 7/SRS.

**Context 1012** [cleaning of previous trench]

### **Samples 9 and 15**

Moist, mid orange-brown, crumbly to stiff (working soft), slightly sandy clay silt. Orange and white

speckles were noted. Stones were present from 6-60+mm in size. Brick/tile, mortar/plaster and charcoal were also present.

These were SRS and BS samples processed on site, no further action was taken as the context was a cleaning layer.

**Context 1037** [machine excavated, mostly medieval and Anglo-Scandinavian finds]

**Samples 28, 29 and 30**

Moist, mid-dark brown (with 1-10mm scale yellowish mottles), soft to sticky (working plastic), sandy silty clay. 2-6mm and 20-60mm stones were present. Other inclusions noted were mortar/plaster, brick/tile, wood, twigs and large mammal bone fragments.

These were SRS and BS samples processed on site; no further action was taken as the context was machine dug. A spot find of hazel nutshell, some specimens with apical knife marks, were noted from Sample 29/SRS.

*Hand-collected shell*

A small quantity of hand-collected oyster shell was recovered from the deposits. The material from the three well-dated contexts is summarised in Table 10.

*Vertebrate remains*

Vertebrate material was recovered from a total of 15 contexts (12 by hand-collection). Material (amounting to ½ box) was recorded from ten of these. The bones from the remaining five (amounting to ½ box) were not recorded, as the contexts were described as either topsoil, back-fill of the previous trench, or machine-excavated deposits. Hand-collected vertebrate material was recorded from seven contexts and bone recovered from four SRS, five BS and 5 GBA samples was also recorded (some contexts having multiple samples).

Preservation records were made for material from three hand-collected contexts and five samples (sometimes from the same contexts). Overall preservation was good, although a few fragments were noted as fair. Colour was slightly variable, being described as brown and dark brown, but the material did not appear mixed in terms of colour. However, the angularity did indicate a degree of reworking of material with battered and rounded fragments present within most contexts.

The degree of fragmentation of the hand-collected material was moderate with most fragments falling between 5 and 20cm in largest dimension. However, the whole assemblage contained a large proportion of small (less than 5cm) fragments. Dog gnawing was noted on less than 10% of the material in two contexts (1013, 1014), whilst fresh breakage was noted in similar quantities but on material in most contexts and samples. Evidence of butchery was noted on up to 20% of the fragments in four contexts.

A total of 1164 fragments (weighing 3024 g) was recorded, of which 104 (weighing 879 g) were identifiable to species or species group. Table 11 shows the numbers of fragments by species together with numbers of measurable and unfused bones, mandibles and loose teeth (of use for ageing or sexing) and weights.

Mammalian species present included the remains of domesticated cattle, caprovid, pig and cat, together with commensal/wild species such as mouse (*Murine* sp.), bank vole (*Clethrionomys glareolus*) and

vole/mouse. Bird species were limited to domestic chicken and a duck which (on the basis of size) could be either wild or domestic. Most of the fish species identified were marine varieties, including herring (*Clupea harengus*), cod (*Gadus morhua*), haddock (*Melanogrammus aeglefinus*) and gadid. Eel (*Anguilla anguilla*) was the only non-marine species present.

The unidentified fraction (1060 fragments) consisted chiefly of completely unidentifiable fragments (688), with large mammal, medium-sized mammal, bird and fish also represented.

Table 12 gives the numbers of fragments by phase and shows that about three-quarters of the fragments were recovered from Phase 4 (all from one context). The numbers of fragments are too small for any differences between phases to be significant. Table 13 shows the numbers of fragments recovered by different techniques. The results are much as would be expected, with the larger more robust fragments recovered by hand-collection and most of the small (particularly fish) bones coming from the fine sieving (BS and GBA samples).

A preliminary examination of the skeletal element representation for the main domesticates (Table 14) indicates that most of the fragments from this site were from non meat-bearing elements (i.e. head and lower limb) for all three species. However, it should be born in mind that the number of fragments was very small and the trends observed may not be significant.

Of the 104 identified fragments, nine were measurable, two had unfused epiphyses, and two mandibles and three isolated teeth (yielding ageing or sexing information) were present.

## **Discussion and statement of potential**

### *Sediment samples*

Plant remains from these samples were generally rather well preserved, either by charring or waterlogging, and in some cases (and unusually) specimens which were only partly charred were recorded. More than one assemblage gave evidence of remains which seemed to have entered the deposit dry and never properly to have become wetted and these, and some material with a sooty 'crust' may indicate the presence of detritus from old thatch. The rye rachis material from at least two contexts was often superbly preserved and occasionally in unbroken lengths of 1-3 cm, an unusual find for deposits in York.

The presence of dyeplant remains, especially of the clubmoss *Diphasium complanatum*, is entirely consistent with the Anglo-Scandinavian date of most of the deposits (and its presence in a context with late medieval pottery warns us against forgetting that it is quite durable and therefore likely to be reworked; the reworking of organic remains preserved by anoxic waterlogging is discussed by Dobney *et al*, 1997). Though never abundant, the remains seem rather unlikely to have reached these deposits by long-distance transport from the huge concentrations seen at 16-22 Coppergate and suggest that dyeing was also practised in the vicinity of Walmgate, though not necessarily at this particular site.

Also characteristic of Anglo-Scandinavian occupation deposits are hazel nuts exhibiting apical knife marks; they were recorded here from Contexts 1004 (Sample 12, 13 and 14), 1005 (8), 1014 (19) and 1037 (29).

No concentrations of intestinal parasitic nematode eggs were encountered. The single observed *Trichuris* egg (Context 1030) was well-preserved, however, implying that none of the examined deposits had a high content of faecal material (rather than that the evidence had not survived).

Preservation of insect remains was much as was observed in Anglo-Scandinavian deposits at 16-22

Coppergate (Kenward and Hall 1995) and Pavement (Hall *et al.* 1983), although perhaps more variable and sometimes showing a considerable degree of erosion or fragmentation (Table 7). Sometimes preservation varied continuously, but in other cases part of the fauna appeared to be in a different preservational state from the majority, and may have had a different origin (e.g. the subjectively recognised heath or moor component in Sample 31, Context 1034). The frequency of charred insects in an identifiable state in one of the samples (Context 1003, Sample 11/T) was exceptional, but charred remains were noted in some of the other samples too.

The overall character of the insect fauna was very similar to material from Coppergate and Pavement. The range of common beetles was indistinguishable, and even the rarer taxa were generally repeatedly observed at Coppergate. ‘House fauna’ was often abundant (Table 8), reflecting an association of most of the Walmgate deposits with buildings. The interiors seems to have been acceptably clean, with indications that conditions became rather moist here and there but probably never really foul. Human fleas were consistently present, sometimes in moderately large numbers, continuing the pattern seen at numerous other sites. Sheep keds, both adults and puparia, and presumably shed by wool cleaning, were present in most of the samples, but not abundant. An aquatic component was generally present (beetles, occasional water fleas of three kinds, and the bryozoan *Lophopus cristallinus*), but no more than might have arrived as background fauna (the site is close to the River Foss) or in water brought to the site.

A number of notable records were made from this material, including the ground beetles *Sphodrus leucophthalmus* (a rarely recorded strongly synanthropic form) and *Odacantha melanura* (found in reed beds and outside its present range at York, Lindroth 1974), the ‘thread-legged bug’ *Empicoris culiciformis* (a genus very rarely recorded from archaeological deposits but likely to have lived in thatch and birds’ nests in the Anglo-Scandinavian town), the delicately patterned froghopper *Paramesus nervosus* (provisionally identified through lack of reference material, a marshland species just outside its present range (Le Quesne 1969, 77), perhaps from the same source as *O. melanura*), and the chafer *Hoplia philanthus*.

### *Shell*

The shell assemblage is too small to be of interpretative value beyond noting that the damage from a knife (or similar implement) to some of the oyster valves indicates the consumption of these shellfish by humans.

### *Vertebrate remains*

The vertebrate assemblage from 41-49 Walmgate was quite small and as such is of limited interpretative value. The preservation was generally good, although the variable angularity of fragments indicates the presence of reworked material. The range of vertebrate species present (particularly fish and small mammals), and the number of small artefacts recovered, has been significantly increased by the extensive sieving programme employed on this site. This shows the value of sieving, even on a small scale excavation, for highlighting the biological and artefactual potential of deposits.

Whilst the recovery of small bones and good preservation of vertebrate material from this excavation has again highlighted the potential deposits in this area of York have for yielding an interesting and useful vertebrate assemblage, the scale of the assemblage recovered is too small for any further work to be undertaken.

## **Recommendations**

No further investigation of the microfossil content of these deposits is recommended.

A proper record of the plant remains from at least some of the samples from this site is required, not least to permit comparison with material of Anglo-Scandinavian date from other sites in York: perhaps Contexts 1003, 1034 and 1039 are the most important. Provision should be made for the preparation of a photographic record of some of the charred rye material, given that it is unusual and extremely well preserved.

The insect remains are clearly important both for reconstructing conditions at the site and in wider synthesis. It would be useful to record remains from larger subsamples in order to make more valid comparisons with sites such as Coppergate and Lloyds Bank.

No further work is recommended on the shell assemblage.

Whilst the vertebrate assemblage is too small to warrant further analytical work, it should be recorded to archive level.

### **Retention and disposal**

All the existing material should be retained and the samples of unprocessed sediment stored under suitable conditions to prevent decay.

### **Archive**

All material is currently stored in the Environmental Archaeology Unit, University of York, along with paper and electronic records pertaining to the work described here.

### **Acknowledgements**

The authors are grateful to Patrick Ottaway (YAT) and Channel 4's 'Time Team' for supplying the material and archaeological information.

### **References**

Carrott, J., Hall, A. And Kenward H. (1991). Environmental evidence from 41-49 Walmgate/George St. (Sitecode: 90.26). *Reports from the Environmental Archaeology Unit, York* **1991/11**.

Carrott, J. and Kenward, H. (2000). Publication draft: Species associations among insect remains from urban archaeological deposits and their significance in reconstructing the past human environment. *Reports from the Environmental Archaeology Unit, York* **2000/11**.

Dainton, M. (1992). A quick, semi-quantitative method for recording nematode gut parasite eggs from archaeological deposits. *Circaea, the Journal of the Association for Environmental Archaeology* **9**, 58-63.

Dobney, K., Hall, A. R., Kenward, H. K. and Milles, A. (1992). A working classification of sample types for environmental archaeology. *Circaea, the Journal of the Association for Environmental Archaeology* **9** (for 1991), 24-6.



- Dobney, K., Kenward, H. and Roskams, S. (1997). All mixed up but somewhere to go? Confronting residuality in bioarchaeology, pp. 81-88 in De Boe, G. and Ver Haeghe, F. (eds.), *Method and theory in historical archaeology*. Papers of the 'Medieval Europe Brugge 1997' Conference **10**. I. A. P. *Rapporten* **10**. Zellik.
- Fisher, R. A., Corbet, A. S. and Williams, C. B. (1943). The relation between the number of species and the number of individuals in a random sample of an animal population. *Journal of Animal Ecology* **12**, 42-58.
- Hall, A. R., Kenward, H. K., Williams, D. and Greig, J. R. A. (1983). Environment and living conditions at two Anglo-Scandinavian sites. *The Archaeology of York* **14** (4), 157-240 plus Plate I and Fiche 1. London: Council for British Archaeology.
- Hall, A. R. (in prep). [Plant remains from Flixborough, N. Lincolnshire.]
- Kenward, H. K. (1992). Rapid recording of archaeological insect remains - a reconsideration. *Circaea, the Journal of the Association for Environmental Archaeology* **9** (for 1991), 81-8.
- Kenward, H. K., Engleman, C., Robertson, A. and Large, F. (1986). Rapid scanning of urban archaeological deposits for insect remains. *Circaea* **3** (for 1985), 163-72.
- Kenward, H. K. and Hall, A. R. (1995). Biological evidence from Anglo-Scandinavian deposits at 16-22 Coppergate. *The Archaeology of York* **14** (7), 435-797 + xxii + loose figures. York: Council for British Archaeology.
- Kenward, H. K., Hall, A. R. and Jones, A. K. G. (1980). A tested set of techniques for the extraction of plant and animal macrofossils from waterlogged archaeological deposits. *Science and Archaeology* **22**, 3-15.
- Kenward, H. and Large, F. (1998). Recording the preservational condition of archaeological insect fossils. *Environmental Archaeology* **2**, 49-60.
- Kloet, G. S. and Hincks, W. D. (1964-77). *A check list of British Insects*. (2nd ed.) London: Royal Entomological Society.
- Le Quesne, W. J. (1969). Hemiptera Cicadomorpha Deltocephalinae. *Handbooks for the identification of British insects* **2** (2b). London: Royal Entomological Society.
- Lindroth, C. H. (1974). Coleoptera Carabidae. *Handbooks for the identification of British insects* **4** (2). London: Royal Entomological Society.
- Peacock, E. R. (1993). Adults and larvae of hide, larder and carpet beetles and their relatives (Coleoptera: Dermestidae) and of derodontid beetles (Coleoptera: Derodontidae). *Handbooks for the identification of British insects* **5** (3). London: Royal Entomological Society.
- Smith, A. J. E. (1978). *The moss flora of Britain and Ireland*. Cambridge: University Press.
- Tutin, T. G. *et al.* (1964-80). *Flora Europaea* **1-5**. Cambridge: University Press.

Table 1. List of samples from 41-49 Walmgate, York with notes on their treatment. Ph-phase

Sample no.	Context	Ph	Approx. sample volume (l.)	Processed? (where?)	Notes	Retain?
1	1000	6	450	Y (On-site)	SRS (10mm mesh)	N/A
2	1000	6	8	Y (On-site)	BS (1mm mesh and washover)	N/A
3	1000	6	60	Y (On-site)	SRS (10mm mesh)	N/A
4	1001	6	120	Y (On-site)	SRS (10mm mesh)	N/A
5	1000	6	15	Y (On-site)	BS (1mm mesh and washover)	N/A
6	1005	1	15	Y (On-site)	SRS (10mm mesh)	N/A
7	1011	6	330	Y (On-site)	SRS (10mm mesh)	N/A
8	1005	1	15	Y (On-site)	BS (1mm mesh and washover)	N/A
9	1012	6	330	Y (On-site)	SRS (10mm mesh)	N/A
10	1005	1	20	Y (EAU)	GBA 1kg subsample, sieved to 300µm with paraffin flotation. 'Squash'.	Y
11	1003	1	10	Y (EAU)	GBA 2kg subsample, sieved to 300µm with paraffin flotation. 'Squash'.	Y
12	1004	1	10	Y (EAU)	GBA 1kg subsample, sieved to 300µm with paraffin flotation. 'Squash'.	Y
13	1004	1	10	Y (On-site)	BS (1mm mesh and washover)	N/A
14	1004	1	10	Y (On-site)	BS (1mm mesh and washover)	N/A
15	1012	6	15	Y (On-site)	BS (1mm mesh and washover)	N/A
16	1013	4	495	Y (On-site)	SRS (10mm mesh)	N/A
17	1011	6	15	Y (On-site)	BS (1mm mesh and washover)	N/A
19	1014	1	10	Y (On-site)	BS (1mm mesh and washover)	N/A
20	1014	1	10	Y (EAU)	GBA 1kg subsample, sieved to 300µm with paraffin flotation. 'Squash'.	Y
21	1014	1	10	Y (On-site)	SRS (10mm mesh)	N/A
22	1030	4	15	Y (On-site)	BS (1mm mesh and washover)	N/A
23	1030	4	210	Y (On-site)	SRS (10mm mesh)	N/A

Sample no.	Context	Ph	Approx. sample volume (l.)	Processed? (where?)	Notes	Retain?
24	1030	4	10	Y (EAU)	GBA 1kg subsample, sieved to 300µm with paraffin flotation. 'Squash'.	Y
25	1030	4	(2 bags)	SPOT		Y
26	1031	3	30	Y (On-site)	BS (1mm mesh and washover)	N/A
27	1031	3	90	Y (On-site)	SRS (10mm mesh)	N/A
28	1037	6	15	Y (On-site)	BS (1mm mesh and washover)	N/A
29	1037	6	420	Y (On-site)	SRS (10mm mesh)	N/A
30	1037	6	10	N		N
31	1034	1	10	Y (EAU)	GBA 1kg subsample, sieved to 300µm with paraffin flotation. 'Squash'.	Y
32	1035	1	10	Y (EAU)	GBA 1kg subsample, sieved to 300µm with paraffin flotation. 'Squash'.	Y
33	1039	1	10	Y (EAU)	GBA 1kg subsample, sieved to 300µm with paraffin flotation. 'Squash'.	Y
34	1037	6	10	N		N

Table 2. Complete list of plant and animal remains recorded from samples from the 41-49 Walmgate, York (Time Team) site, in taxonomic order. Although one sample came from a supposedly later medieval deposit, its flora and fauna appeared to be of Anglo-Scandinavian date and the records for it are included here. Order and nomenclature follow Tutin et al. (1964-90) for vascular plants, Smith (1976) for mosses, and Kloet and Hincks (1964-77) for insects. Plant material not specifically noted as being preserved by charring can be taken to be uncharred. Where both secure and tentative identifications for a given taxon were recorded, only the former are listed here. For invertebrates, \* = not used in calculating assemblage statistics (Table 8); ecode - ecological code used in generating main statistics (Tables 4 and 6); Sp(p). - species not previously listed; Sp(p). indet. - may be a species already listed.

BRYOPHYTA (all remains were leaves and/or shoots unless otherwise specified)

*Sphagnum* sp(p). leaves  
*Ceratodon purpureus* (Hedw.) Brid.  
*Dicranum scoparium* Hedw.  
*Leucobryum glaucum* (Hedw.) Ångstr.  
*Barbula* sp(p).  
*Racomitrium canescens* (Hedw.) Brid.  
*Ulota* sp(p).  
*Leucodon sciuroides* (Hedw.) Sch waegr.  
*Antitrichia curtipendula* (Hedw.) Brid.  
*Neckera complanata* (Hedw.) Hüb.  
*Thuidium tamariscinum* (Hedw.) Br. Eur.  
 cf. *Cratoneuron filicinum* (Hedw.) Spruce  
*Campylium stellatum* (Hedw.) Lange & Jens.  
*C. cf. elodes* (Lindb.) Kindb.  
 cf. *Amblystegium riparium* (Hedw.) Br. Eur.  
*Drepanocladus* sp(p).  
*Scorpidium scorpioides* (Hedw.) Limpr.  
*Calliergon* cf. *giganteum* (Schimp.) Kindb.  
*C. cuspidatum* (Hedw.) Kindb.  
*Isoetecium myurum* Brid.  
*I. myosuroides* Brid.  
*Homalothecium sericeum* (Hedw.) Br. Eur./*H. lutescens* (Hedw.) Robins.  
*Eurhynchium striatum* (Hedw.) Schimp.  
*Eurhynchium* sp(p).  
*Hypnum* cf. *cupressiforme* Hedw.  
*Rhytidiadelphus* cf. *squarrosus* (Hedw.) Warnst.  
*Rhytidiadelphus* sp(p).  
*Hylocomium* cf. *brevirostre* (Brid.) Br. Eur.  
*H. splendens* (Hedw.) Br. Eur.

PTERIDOPHYTA

*Diplazium complanatum* (L.) Rothm. (complanate clubmoss) charred and uncharred shoot fragments  
*Pteridium aquilinum* (L.) Kuhn (bracken) pinnae fragments

ANGIOSPERMAE

*Salix* sp(p). (willow) buds  
*Populus* sp(p). (poplar/aspens) buds and/or bud-scales  
*Myrica gale* L. (bog myrtle/sweet gale) leaf fragments  
*Juglans regia* L. (walnut) nutshell fragments  
*Betula* cf. *pendula* Roth (?silver birch) bark fragments  
*Betula* sp(p). (birch) female catkin scales, fruits  
*Alnus glutinosa* (L.) Gaertner (alder) buds and/or bud-scales, fruits  
*Corylus avellana* L. (hazel) charred and uncharred nuts and/or nutshell fragments, immature nuts  
*Quercus* sp(p). (oak) buds and/or bud-scales, wood fragments  
*Ficus carica* L. (fig) seeds  
*Humulus lupulus* L. (hop) achenes  
*Cannabis sativa* L. (hemp) achenes  
*Urtica dioica* L. (stinging nettle) achenes  
*U. urens* L. (annual nettle) achenes  
*Polygonum aviculare* agg. (knotgrass) charred and uncharred fruits  
*P. hydropiper* L. (water-pepper) fruits  
*P. persicaria* L. (persicaria/red shank) fruits  
*P. persicaria/lapathifolium* (persicarias) charred and uncharred fruits  
*P. lapathifolium* L. (pale persicaria) fruits  
*Bilderdykia convolvulus* (L.) Dumort. (black bindweed) charred and uncharred fruits and uncharred fruit fragments  
*Rumex acetosella* agg. (sheep's sorrel) fruits  
*Rumex* sp(p). (docks) charred and uncharred fruits, perianth s/perianth segments,  
 cf. *Rumex* sp(p). (?docks) flowering stem fragments  
*Chenopodium* Section *Pseudoblitum* (red goosefoot, etc.) seeds  
*C. murale* L. (nettle-leaved goosefoot) seeds  
*C. album* L. (fat hen) charred and uncharred seeds  
*Atriplex* sp(p). (oraches) charred and uncharred seeds  
*Stellaria media* (L.) Vill. (chickweed) seeds  
*S. palustris* Retz./*S. graminea* L. (marsh/lesser stitchwort) seeds  
*Spergula arvensis* L. (com spurrey) seeds

*Agrostemma githago* L. (comcockle) charred and uncharred seeds and uncharred seed fragments  
*Silene vulgaris* (Moench) Garcke (bladder campion) seeds  
*S. alba* (Miller) Krause in Sturm (white campion) seeds  
*Caltha palustris* L. (marsh marigold) seeds  
*Ranunculus* Section *Ranunculus* (meadow/creeping/bulbous buttercup) achenes  
*R. sardous* Crantz (hairy buttercup) achenes  
*R. flammula* L. (lesser spearwort) achenes  
 Cruciferae (cabbage family) pedicels, seeds  
*Isatis tinctoria* L. (woad) pod fragments  
*Camelina sativa* (L.) Crantz (gold-of-pleasure) pods  
*Thlaspi arvense* L. (field penny-cress) seeds  
*Brassica rapa* L. (tumpip) seeds  
*Brassica* sp(p). (cabbages, etc.) seeds  
*Brassica* sp./*Sinapis arvensis* (brassica/charlock) mineralised cotyledons and uncharred seeds  
*Raphanus raphanistrum* L. (wild radish) charred and uncharred pod segments and/or fragments and uncharred seeds  
*Reseda luteola* L. (weld/dyer's rocket) seeds  
*Filipendula ulmaria* (L.) Maxim. (meadowsweet) achenes  
*Rubus fruticosus* agg. (blackberry/bramble) seeds  
*Rubus/Rosa* sp(p). (blackberry, etc./rose) prickles  
*Rosa* sp(p). (roses) achenes  
*Potentilla palustris* (L.) Scop. (marsh cinquefoil) achenes  
*P. cf. erecta* (L.) Räuschel (?tormentil) achenes  
*Fragaria cf. vesca* L. (?wild strawberry) charred and uncharred achenes  
*Aphanes arvensis* agg. (parsley-pierts) achenes  
*Malus sylvestris* Miller (crab apple) endocarp, seeds and seed fragments  
*Prunus spinosa* L. (sloe) charred and uncharred fruitstones, thorns  
*P. domestica* ssp. *insititia* (L.) C. K. Schneider (plums, etc.) fruitstones  
 Leguminosae (pea family) calyx/calyces, flowers and/or petals, immature seeds (waterlogged), pods and/or pod fragments  
*Genista tinctoria* L. (dyer's greenweed) stem fragments, twig epidermis fragments  
*Vicia faba* L. (field bean) waterlogged hila  
 cf. *Vicia* sp(p). (?vetches, etc.) charred seeds  
*Pisum cf. sativum* L. (?garden/field pea) charred seeds  
*Oxalis acetosella* L. (wood-sorrel) seeds  
*Linum usitatissimum* L. (cultivated flax) capsule fragments, charred and uncharred seeds  
*Euphorbia helioscopia* L. (sun spurge) seeds  
*Viola* sp(p). (violets/pansies, etc.) capsule segments, charred and uncharred seeds  
 Umbelliferae (carrot family) mericarps  
*Pimpinella saxifraga* L. (burnet saxifrage) mericarps  
*Oenanthe cf. aquatica* (L.) Poiret in Lam. (?fine-leaved water-dropwort) mericarps  
*Aethusa cynapium* L. (fool's parsley) charred and uncharred mericarps  
*Anethum graveolens* L. (dill) mericarps  
*Torilis japonica* (Houtt.) DC. (upright hedge-parsley) mericarps  
*Daucus carota* L. (wild carrot) mericarps  
*Erica tetralix* L. (cross-leaved heath) leaf/leaves  
*Calluna vulgaris* (L.) Hull (heather, ling) shoot fragments  
 cf. *C. vulgaris* (?heather, ling) charred and uncharred root and/or basal twig fragments  
 Primulaceae (primrose family) seeds  
*Anagallis arvensis* L. (scarlet pimpernel) seeds  
*Menyanthes trifoliata* L. (bogbean) seeds  
*Rubia tinctorum* L. (dyer's madder) root bark and root fragments  
*Marrubium vulgare* L. (white horehound) nutlets  
*Galeopsis* Subgenus *Ladanum* (hemp-nettles) nutlets  
*G.* Subgenus *Galeopsis* (hemp-nettles) nutlets  
*Lamium* Section *Lamiopsis* (annual dead-nettles) nutlets  
*Stachys* sp(p). (woundworts) nutlets  
*Nepeta cataria* L. (cat-mint) nutlets  
*Prunella vulgaris* L. (selfheal) nutlets  
*Satureja hortensis* L. (summer savory) nutlets  
*Hyoscyamus niger* L. (henbane) seeds  
*Pedicularis palustris* L. (marsh lousewort) seeds  
*Plantago major* L. (greater plantain) seeds  
*Sambucus nigra* L. (elder) charred and uncharred seeds  
*Valerianella dentata* (L.) Pollich (narrow-fruited cornsalad) fruits  
*Knautia arvensis* (L.) Coulter (field scabious) fruit fragments  
 Compositae (daisy family) involucre/fragments  
*Bellis perennis* L. (daisy) achenes  
*Aster tripolium* L. (sea aster) achenes  
*Bidens* sp(p). (bur-marigolds) achenes  
*Anthemis cotula* L. (stinking mayweed) charred and uncharred achenes  
*Arctium* sp(p). (burdocks) achenes  
*Carduus/Cirsium* sp(p). (thistles) achenes  
*Centaurea* sp(p). (knapweeds, etc.) achenes  
*Hypochoeris* sp(p). (cat's ears) achenes  
*Leontodon* sp(p). (hawkbits) achenes  
*Picris hieracioides* L. (hawkweed ox-tongue) achenes  
*Sonchus asper* (L.) Hill (prickly sow-thistle) achenes  
*S. oleraceus* L. (sow-thistle) achenes  
*S. palustris* L./*S. arvensis* L. (marsh/corn sow-thistle) achenes

<i>S. cf. arvensis</i> L. (?corn sow-thistle) achenes		* <i>Daphnia</i> sp. (ephippium)	oa-w
<i>Lapsana communis</i> L. (nipplewort) achenes		* <i>Cladocera</i> sp. (ephippium)	oa-w
<i>Triglochin maritima</i> L. (sea arrowgrass) charred and uncharred carpels			
<i>Iris pseudacorus</i> L. (yellow flag) seeds		INSECTA	
<i>Juncus cf. compressus</i> Jacq. (?round-fruited rush) seeds		DERMAPTERA	
<i>J. cf. gerardi</i> Loisel. (?mud rush) seeds		* <i>Forficula auricularia</i> Linnaeus	rt
<i>J. bufonius</i> L. (toad rush) seeds		HEMIPTERA	
<i>Luzula</i> sp(p). (woodrushes) seeds		<i>Empicoris culiciformis</i> (Degeer)	rt
Gramineae (grasses) charred and uncharred caryopsis/es		<i>Lyctocoris campestris</i> (Fabricius)	rd-st
Gramineae/'Cerealia' (grasses/cereals) charred culm fragments, culm nodes, awns/awn fragments, caryopses, and rachis fragments and uncharred culm nodes		<i>Chartoscirta ? cocksi</i> (Curtis)	oa-w
cf. <i>Glyceria</i> sp(p). (?sweet-grasses) caryopses		* <i>Heteroptera</i> sp. (nymph)	u
<i>Bromus</i> sp(p). (bromes, etc.) charred caryopses		<i>Aphrodes</i> sp.	oa-p
<i>Triticum 'aestivo-compactum'</i> (bread/club wheat) charred caryopsis/es		? <i>Paramesus nervosus</i> (Fallen)	oa-p
<i>Triticum</i> sp(p). (wheats) charred caryopses		Delphacidae spp.	oa-p
<i>Triticum/Secale</i> (wheat/rye) waterlogged caryopses and periderm ('bran') fragments		Auchenorrhyncha spp.	oa-p
<i>Secale cereale</i> L. (rye) charred caryopses, and lemma and rachis fragments		* <i>Auchenorrhyncha</i> sp. (nymph)	oa-p
cf. <i>Secale cereale</i> L. (?rye) charred caryopses and uncharred rachis fragments		* <i>Coccoidea</i> sp.	u
<i>Hordeum vulgare</i> L. (six-row barley) charred caryopsis/es, some twisted grains		? <i>Hemiptera</i> sp.	u
<i>Hordeum</i> sp(p). (barley) charred caryopses		TRICHOPTERA	
<i>Avena fatua</i> L. (wild oat) charred spikelets/spikelet fragments		*?Trichoptera sp.	oa-w
<i>A. sativa</i> L. (cultivated oat) charred and uncharred spikelets/spikelet fragments		LEPIDOPTERA	
<i>Avena</i> sp(p). (oats) charred caryopses, chaff, part-charred spikelets/spikelet fragments, uncharred caryopses and spikelets/spikelet fragments		* <i>Lepidoptera</i> sp. (pupa)	u
<i>Danthonia decumbens</i> (L.) DC. in Lam. & DC. (heath grass) caryopses		DIPTERA	
<i>Scirpus maritimus</i> L. (sea club-rush) nutlets		* <i>Melophagus ovinus</i> (Linnaeus) (puparium)	u
<i>S. setaceus</i> L. (bristle club-rush) nutlets		* <i>Melophagus ovinus</i> (Linnaeus) (adult)	oa-w
<i>Eleocharis palustris</i> s.l. (common spike-rush) charred and uncharred nutlets		* <i>Diptera</i> sp. (adult)	u
<i>Eleocharis</i> sp(p). nutlets		* <i>Diptera</i> sp. (larva)	u
<i>Cladium mariscus</i> (L.) Pohl (great sedge/saw-sedge) nutlets with perigynium		* <i>Diptera</i> sp. (puparium)	u
cf. <i>C. mariscus</i> (L.) Pohl (?great sedge/saw-sedge) charred leaf fragments		* <i>Diptera</i> sp. (pupa)	u
<i>Schoenus nigricans</i> L. (bog-rush) nutlets		* <i>Syrphidae</i> sp. (larva)	u
<i>Carex</i> sp(p). (sedges) charred and uncharred nutlets		SIPHONAPTERA	
	e-code	* <i>Pulex irritans</i> Linnaeus	ss
ANNELIDA: OLIGOCHAETA		* <i>Siphonaptera</i> sp. indet.	u
* <i>Oligochaeta</i> sp. (egg capsule)	u	COLEOPTERA	
CRUSTACEA: CLADOCERA		? <i>Sphodrus leucophthalmus</i> (Linnaeus)	oa
		<i>Clivina fossor</i> (Linnaeus)	oa
		<i>Trechus quadristriatus</i> (Schrank)	oa
		<i>Trechus micros</i> (Herbst)	u
		? <i>Trechus</i> sp.	ob
		<i>Bembidion (Philochthus)</i> sp.	oa
		<i>Bembidion</i> sp.	oa
		<i>Stomis pumicatus</i> (Panzer)	oa
		<i>Pterostichus (Poecilus)</i> sp.	oa
		<i>Pterostichus</i> sp.	ob
		<i>Agonum dorsale</i> (Pontoppidan)	oa
		<i>Amara</i> sp.	oa
		<i>Odacantha melanura</i> (Linnaeus)	oa-d
		Carabidae spp. and spp. indet.	ob
		<i>Helophorus ?aquaticus</i> (Linnaeus)	oa-w
		<i>Helophorus aquaticus</i> or <i>grandis</i>	oa-w
		<i>Helophorus</i> spp.	oa-w

<i>Cercyon analis</i> (Paykull)	rt-sf	<i>Crataraea suturalis</i> (Mannerheim)	rt-st
<i>Cercyon haemorrhoidalis</i> (Fabricius)	rf-sf	Aleocharinae spp.	u
<i>Cercyon unipunctatus</i> (Linnaeus)	rf-st	Staphylinidae sp.	u
<i>Cryptopleurum minutum</i> (Fabricius)	rf-st	Euplectini sp.	u
<i>Acritus nigricornis</i> (Hoffmann)	rt-st	Pselaphidae sp.	u
Histerinae sp.	rt	<i>Trox scaber</i> (Linnaeus)	rt-sf
<i>Ochthebius ? minimus</i> (Fabricius)	oa-w	<i>Aphodius granarius</i> (Linnaeus)	ob-rf
<i>Ochthebius</i> sp. indet.	oa-w	<i>Aphodius prodromus</i> (Brahm)	ob-rf
<i>Hydraena</i> sp.	oa-w	<i>Aphodius</i> spp. and spp. indet.	ob-rf
<i>Ptenidium</i> sp.	rt	<i>Hoplia philanthus</i> Illiger	oa
<i>Acrotrichis</i> sp.	rt	<i>Clambus ? pubescens</i> Redtenbacher	rt-sf
<i>Catops</i> sp.	u	<i>Clambus</i> sp.	rt-sf
<i>Silpha atrata</i> Linnaeus	u	<i>Cyphon</i> sp.	oa-d
<i>Scydmaenus</i> sp.	u	Elatenidae spp.	ob
<i>Micropeplus fulvus</i> Erichson	rt	* <i>Dermestes lardarius</i> (larva)	rd-sf
<i>Micropeplus</i> sp. indet.	rt	<i>Anobium punctatum</i> (Degeer)	l-sf
<i>Phyllodrepa ? floralis</i> (Paykull)	rt-sf	<i>Ptinus ? fur</i> (Linnaeus)	rd-sf
<i>?Phyllodrepa salicis</i> (Gyllenhal)	rt-sf	<i>Ptinus</i> sp. indet.	rd-sf
<i>Dropephylla ? ioptera</i> (Stephens)	u	Ptinidae sp. indet.	rd
<i>Dropephylla vilis</i> (Erichson)	l	<i>Lyctus linearis</i> (Goeze)	l-sf
<i>Dropephylla</i> sp. indet.	u	<i>Meligethes</i> sp.	oa-p
<i>Omalium excavatum</i> Stephens	rt-sf	<i>Omosita ? discoidea</i> (Fabricius)	rt-sf
<i>Omalium caesum</i> or <i>italicum</i>	rt-sf	<i>Omosita</i> sp. indet.	rt-sf
<i>Omalium ? rivulare</i> (Paykull)	rt-sf	<i>Rhizophagus</i> sp.	u
<i>Omalium</i> sp. indet.	rt	<i>Monotoma spinicollis</i> Aube	rt-st
<i>Xylodromus concinnus</i> (Marsham)	rt-st	<i>Monotoma</i> sp. indet.	rt-sf
<i>Coprophilus striatulus</i> (Fabricius)	rt-st	<i>Cryptophagus scutellatus</i> Newman	rd-st
<i>Carpelimus bilineatus</i> Stephens	rt-sf	<i>Cryptophagus</i> spp.	rd-sf
<i>Carpelimus fuliginosus</i> (Gravenhorst)	st	<i>Atomaria nigripennis</i> (Kugelann)	rd-ss
<i>Carpelimus pusillus</i> group	u	<i>Atomaria</i> spp.	rd
<i>Carpelimus</i> sp.	u	<i>Ephistemus globulus</i> (Paykull)	rd-sf
<i>Platystethus arenarius</i> (Fou rcroy)	rf	<i>Orthoperus</i> sp.	rt
<i>Platystethus degener</i> Mulsant & Rey	oa-d	<i>Mycetaea hirta</i> (Marsham)	rd-ss
<i>Platystethus cornutus</i> group indet.	oa-d	<i>Lathridius minutus</i> group	rd-st
<i>Platystethus nitens</i> (Sahlberg)	oa-d	<i>Enicmus</i> sp.	rt-sf
<i>Anotylus complanatus</i> (Erichson)	rt-sf	<i>Corticaria</i> spp.	rt-sf
<i>Anotylus nitidulus</i> (Gravenhorst)	rt	<i>Typhaea stercorea</i> (Linnaeus)	rd-ss
<i>Anotylus rugosus</i> (Fabricius)	rt	<i>Aglenus brunneus</i> (Gyllenhal)	rt-ss
<i>Anotylus sculpturatus</i> group	rt	<i>Tenebrio obscurus</i> Fabricius	rt-ss
<i>Oxytelus sculptus</i> Gravenhorst	rt-st	Salpingidae sp.	l
<i>Stenus</i> spp.	u	<i>Anthicus formicarius</i> (Goeze)	rt-st
<i>Lathrobium</i> sp.	u	<i>Anthicus floralis</i> or <i>formicarius</i>	rt-st
<i>Leptacinus pusillus</i> (Stephens)	rt-st	<i>Phymatodes alni</i> (Linnaeus)	l
<i>Leptacinus</i> sp. indet.	rt-st	Cerambycidae sp.	l
<i>Phacophallus parumpunctatus</i> (Gyllenhal)	rt-st	<i>Phaedon</i> sp.	oa-p
<i>Gyrophypnus angustatus</i> Stephens	rt-st	<i>Phyllotreta nemorum</i> group	oa-p
<i>Gyrophypnus fracticornis</i> (Muller)	rt-st	<i>Phyllotreta</i> sp.	oa-p
<i>Neobisnius</i> sp.	u	<i>Longitarsus</i> sp.	oa-p
<i>Philonthus ? politus</i> (Linnaeus)	rt-st	<i>? Altica</i> sp.	oa-p
<i>Philonthus</i> spp.	u	<i>Chaetocnema ? concinna</i> (Marsham)	oa-p
<i>Creophilus maxillosus</i> (Linnaeus)	rt	<i>Chaetocnema</i> sp.	oa-p
<i>Quedius</i> sp.	u	Halticinae sp.	oa-p
Staphylininae spp. and spp. indet.	u	<i>Apion (Oxystoma) ? subulatum</i> Kirby	oa-p
<i>Tachinus laticollis</i> or <i>marginellus</i>	u	<i>Apion</i> spp.	oa-p
<i>Cordalia obscura</i> (Gravenhorst)	rt-sf	<i>? Strophosomus</i> sp.	oa-p
<i>Falagria caesa</i> or <i>sulcatula</i>	rt-sf	<i>Sitona</i> sp.	oa-p
<i>Falagria</i> or <i>Cordalia</i> sp.	rt-sf	Cossoninae sp.	u

<i>Dorytomus</i> sp.	oa-p
<i>Ceutorhynchus contractus</i> (Marsham)	oa-p
<i>Ceutorhynchus</i> sp.	oa-p
Ceuthorhynchinae sp.	oa-p
Curculionidae sp.	oa
<i>Leperisinus ?varius</i> (Fabricius)	l
*Coleoptera sp. indet. (larva)	u

HYMENOPTERA

*Proctotrupoidea sp.	u
*Chalcidoidea sp.	u
*Hymenoptera Parasitica sp.	u
*Formicidae sp.	u
*? <i>Apis mellifera</i> Linnaeus	u
*Apoidea sp. indet.	u
*Hymenoptera sp.	u
*Insecta sp. (larva)	u

ARACHNIDA

*Aranae sp.	u
*Acarina sp.	u

MOLLUSCA

*Gastropoda sp.	u
-----------------	---

BRYOZOA

* <i>Lophopus crystallinus</i> (Pallas)	oa-w
---	------

VERTEBRATA

PISCES

*Clupea Harengus* L. (herring)  
*Anguilla anguilla* (L.) (eel)  
Gadiidae (cod family)  
*Gadus morhua* L. (cod)  
*Melanogrammus aeglefinus* (L.) (haddock)

AVES

*Anas* sp. (duck)  
cf. *Gallus* f. domestic (?fowl)  
*Gallus* f. domestic (chicken)

MAMMALIA

Microtine/Murine (vole/mouse)  
*Clethrionomys glareolus* (Screber) (bank vole)  
Murinae (mouse species)  
*Felis* f. domestic (cat)  
*Sus* f. domestic (pig)  
*Bos* f. domestic (cattle)  
Caprovid (sheep/goat)



Table 3. Lists of plants remains and other components of the samples from 41-9 Walmgate (Time Team site), York, in context, sample and subsample order. For each list records are presented in descending order by abundance score (on a 3- or 4-point scale as appropriate for the kind of sample) and for each score in alphabetical order.

Abbreviations: *af*—achene fragment(s); *b*—bud(s); *br*—bract(s); *bs*—bud-scale(s); *caps*—capsule(s); *ch*—charred; *c/n*—culm-nodes; *cot*—cotyledon(s); *dec*—decayed; *dim*—dimension; *endo*—endocarp; *epid*—epidermis; *fc*s—female catkin or cone scale(s); *ff*—fruit fragment(s); *fgt/s*—fragment(s); *fls*—flower(s); *fr*—fruit(s); *inc*—including; *incomp*—incompletely; *inv*—involucre/involucral; *lef*—leaf epidermis fragment(s); *lvs*—leaves; *max*—maximum; *mc*—male catkin; *meso*—mesocarp; *mf*—mericarp fragment(s); *min*—mineral-replaced ('mineralised'); *per*—perianth(s); *pet*—petal(s); *pinn*—pinnule; *rt*—root; *s*—seed(s); *segs*—segment(s); *sf*—seed fragment(s); *sh*t—shoot; *specn*—specimen; *spklts*—spikelet(s); *spr*—sprouting; *st*—stem; *tef*—twig epidermis fragment(s); *tw*—twig; *twstd*—twisted; *v*—very.

Context 1000, Sample 5/BS		earthworm egg caps	1
Diphasium complanatum	1	fish bone	1 max 2mm
		Gramineae (ch)	1
		gravel	1 max 20mm
Context 1003, Sample 11/T		herbaceous detritus	1
charcoal	3 max 40mm	Hordeum sp(p).	1
charred herbaceous detritus	3	Linum usitatissimum (caps fgts)	1
sand	3	Menyanthes trifoliata	1 fragment(s) only
Agrostemma githago (ch)	2	moss (contaminant)	1
Gramineae/Cerealialia (c/n)	2	Neckera complanata	1
Secale cereale (lemma fgts)	2	Polygonum aviculare agg. (ch)	1
Secale cereale (rachis fgts)	2	Polygonum persicaria/lapathifolium (ch)	1
'ash beads'	1	Quercus sp(p). (b/bs)	1
'char'	1 max 10 mm	Raphanus raphanistrum (pod segs/fgts)	1
'pinched' stems (ch)	1	part-charred	(toasted)
?daub	1 max 40mm	Rubus fruticosus agg.	1
Aethusa cynapium (ch)	1	Rumex sp(p). (ch)	1
animal hairs	1	Sambucus nigra	1
Anthemis cotula	1	Sambucus nigra (ch)	1
Anthemis cotula (ch)	1	Scorpidium scorpioides	1
ash concretions	1 max 5mm	Secale cereale	1
Atriplex sp(p).	1	Secale cereale (part-ch rachis fgts)	1
Avena sativa (spklts/fgts)	1	snails	1
Avena sp(p).	1	Sphagnum sp(p). (lvs)	1
bark fgts	1 max 20mm	Triglochin maritima (ch)	1
Bilderdykia convolvulus (ch)	1	Triticum aestivo-compactum	1
Brassica rapa	1	Urtica urens	1
burnt bark fgts	1 max 10 mm	Viola sp(p). (ch)	1
burnt bone fgts	1 max 20mm	wood chips	1 max 10 mm
Calliargon cf. giganteum	1	wood chips (ch)	1 some or all only partly charred; 10 mm max dim
Carex sp(p). (ch)	1		
Centaurea sp(p).	1		
Cerealialia in det. (chaff)	1		
Cerealialia indet. (rachis fgts)	1		
cf. Avena sp(p). (part-ch chaff)	1		
cf. Secale cereale	1		
cf. Vicia sp(p).	1 max 2mm		
charred trigonous stems	1		
Chenopodium album	1	Context 1004, Sample 12/T	
Chenopodium album (ch)	1	Anthemis cotula	2
Chenopodium Section Pseudoblitum	1	Atriplex sp(p).	2
Corylus avellana	1	bark fgts	2 max 25mm
		Brassica rapa	2 inc fgts

charcoal	2 max 15mm	Juncus cf. compressus	1
Chenopodium album	2	Juncus cf. gerardi	1
earthworm egg caps	2	Lapsana communis	1
herbaceous detritus	2 mostly <1mm	Leucobryum glaucum	1
Raphanus raphanistrum (pod segs/fgts)	2	Linum usitatissimum	1 inc fgts
sand	2	Linum usitatissimum (caps fgts)	1
Urtica urens	2	Marrubium vulgare	1
wood chips	2 max 10 mm	Melophagus ovinus (sheep ked)	1
'char'	1 max 5mm	Neckera complanata	1
'coils'	1	Polygonum aviculare agg.	1
?peat/mor humus	1 max 5mm	Polygonum persicaria	1
Aethusa cynapium	1	Potentilla cf. erecta	1
Agrostemma githago	1	pottery	1 max 30mm
Agrostemma githago (sf)	1	Prunella vulgaris	1
Anagallis arvensis	1	Quercus sp(p). (b/bs)	1
Anthemis cotula (ch)	1	Ranunculus sardous	1
Avena sp(p).	1	Raphanus raphanistrum (s)	1 fragment(s)
Avena sp(p). (chaff)	1	only	
Avena sp(p). (part-ch spkls/fgts)	1	root/rootlet fgts	1
Avena sp(p). (w/l spkls/fgts)	1	Rubia tinctorum	1 max 2mm
Avena sp(p). (w/l)	1	Rumex acetosella agg.	1
bark chips	1 max 5mm	Rumex sp(p).	1
beetles	1	Sambucus nigra	1
Bellis perennis	1	Satureja hortensis	1
Bilderdykia convolvulus	1	Scirpus setaceus	1
bone fgts	1 max 10 mm	Scorpidium scorpioides	1
Brassica sp./Sinapis arvensis	1	Sonchus asper	1
brick/tile	1 max 5mm	Sonchus palustris/arvensis	1
burnt bone fgts	1 max 5mm	Spergula arvensis	1
Calliargon cf. giganteum	1	Stellaria media	1
Camelina sativa (pods)	1	Stellaria palustris/graminea	1
Carex sp(p).	1	teeth	1 fragment(s) only
Cenococcum (sclerotia)	1	Valerianella dentata	1
Cerealia indet. (awns)	1	Viola sp(p).	1
Cerealia in det. (chaff)	1	Viola sp(p). (caps segs)	1
cf. Rumex sp(p). (flg st fgts)	1	wood fgts	1 max 15mm
Corylus avellana	1 including material with apical knife marks		
<hr/>			
Context 1004, Sample 13/BS			
Danthonia decumbens	1	Anthemis cotula	2
Diphasium complanatum	1 very	Avena sp(p).	2
decayed		Chenopodium album	2
Eleocharis palustris sl	1	Corylus avellana	2 including material with apical knife marks
fish bone	1 max 10 mm		
fly puparia	1	Lapsana communis	2
Galeopsis Subgenus Galeopsis	1	Linum usitatissimum	2
Gramineae	1	wood fgts	2 max 20mm
Gramineae/Cerealia (ch culm fgts)	1	Aethusa cynapium	1
gravel	1 max 20mm	Agrostemma githago	1
grit	1	Agrostemma githago (sf)	1
Hyoscyamus niger	1	Anethum graveolens	1
Hypnum cf. cupressiforme	1	Atriplex sp(p).	1
Isatis tinctoria (pod fgts)	1	Avena fatua (spkls/fgts)	1
Isoetecium myurum	1	Avena sativa (spkls/fgts)	1
Juncus bufonius	1		

Avena sativa (w/l spkls/fgts)	1		spec: tubed
Avena sp(p). (chaff)	1	Polygonum aviculare agg.	1
bark fgts	1 max 10 mm	Polygonum lapathifolium	1
bast with sclereids	1	Polygonum persicaria	1
beetles	1	pottery	1 max 10 mm
Bilderdykia convolvulus	1	Prunella vulgaris	1
Bilderdykia convolvulus (ff)	1	Racomitrium canescens	1
bone fgts	1 max 20mm	Ranunculus flammula	1
Brassica rapa	1	Ranunculus sardous	1
Brassica sp./Sinapis arvensis	1	Ranunculus Section Ranunculus	1
burnt bone fgts	1 max 15mm	Raphanus raphanistrum (pod segs/fgts)	1
Calliergon cf. giganteum	1	Rumex sp(p).	1
Carex sp(p).	1	Sambucus nigra	1
	nutlets with utricles	sand	1
	and/or free utricles	Scorpidium scorpioides	1
		Secale cereale (lemma fgts)	1 a single spec: tubed
Cerealia in det. (chaff)	1		
Cerealia in det. (w/l chaff)	1	Secale cereale (rachis fgts)	1
cf. Avena sp(p). (chaff)	1	Sonchus asper	1
cf. Calluna vulgaris (ch rt-tw fgts)	1	Spergula arvensis	1
cf. Calluna vulgaris (rt-tw fgts)	1	Sphagnum sp(p). (lvs)	1
cf. Secale cereale	1	Stachys sp(p).	1
cf. Secale cereale (w/l rachis fgts)	1	Stellaria media	1
cf. Triticum/Secale (w/l)	1	Thuidium cf. tamariscinum	1
charcoal	1 max 15mm	Torilis japonica	1
charred herbaceous detritus	1	Triglochin maritima	1
Cladium mariscus (with perig)	1	Triticum aestivo-compactum	1
coal	1 max 10 mm	Umbelliferae	1
Corylus avellana (ch)	1	unwashed organic sediment	1 max 30mm
Corylus avellana (imm)	1	Urtica dioica	1
Diphasium complanatum	1 very	Urtica urens	1
decayed		Valerianella dentata	1
eggshell fgts	1 max 5mm	Viola sp(p).	1
Eleocharis palustris sl	1	wattle/wicker fgts	1 max 25mm
Eurhynchium sp(p).	1	wood chips	1 max 10 mm
fish bone	1 max 20mm		
fish scale	1 max 5mm		
Galeopsis Subgenus Galeopsis	1	<hr/> Context 1004, Sample 14/SPT <hr/>	
Genista tinctoria (tef)	1 a single spec	Corylus avellana	1 including material
gravel	1 max 10 mm		with apical knife marks
Hordeum sp(p).	1		
Hyoscyamus niger	1	Corylus avellana	1
Hypnum cf. cupressiforme	1	fish bone	1 max 15mm
Leguminosae (cal)	1		
Leguminosae (pods/fgts)	1 max 5mm		
Leontodon sp(p).	1	<hr/> Context 1005, Sample 8/BS <hr/>	
Leucobryum glaucum	1	wood fgts	3 max 15mm
Linum usitatissimum (caps fgts)	1	Anthemis cotula	2
Malus sylvestris	1	Atriplex sp(p).	2
Malus sylvestris (endo)	1	bark fgts	2 max 45mm
Malus sylvestris (seed base cups)	1	charcoal	2 max 20mm
mammal tooth	1	Chenopodium album	2
Melophagus ovinus (sheep ked)	1	gravel	2 max 20mm
moss (contaminant)	1	grit	2
Neckera complanata	1	Linum usitatissimum	2
Pimpinella saxifraga	1 a single		

Urtica dioica	2	Humulus lupulus	1
wood chips	2 max 10 mm	Hylocomium splendens	1
?daub	1 max 30mm	Hypochoeris sp(p).	1
Aethusa cynapium	1	Iris pseudacorus	1 fragment(s) only
Arctium sp(p).	1	Isatis tinctoria (pod fgts)	1
Avena cf. sativa (spklt/sfgts)	1	Isothecium myosuroides	1
Avena sp(p).	1	Isothecium myurum	1
Avena sp(p). (chaff)	1	Juncus bufonius	1
beetles	1	Knautia arvensis (ff)	1
Bidens sp(p).	1	Lapsana communis	1
Bilderdykia convolvulus	1	leather fgts	1 max 10 mm
Bilderdykia convolvulus (ff)	1	Leguminosae (cal)	1
bone fgts	1 max 20mm	Leguminosae (fls/pet)	1
Brassica rapa	1	Leontodon sp(p).	1
Brassica sp(p).	1	Leucobryum glaucum	1
Brassica sp./Sinapis arvensis	1	Malus sylvestris	1
brick/tile	1 max 5mm	mammal tooth	1
Bromus sp(p).	1	Melophagus ovinus (sheep ked)	1
burnt bone fgts	1 max 20mm	Menyanthes trifoliata	1
Calliergon cuspidatum	1	moss (contaminant)	1
Calluna vulgaris (sht fgts)	1	Neckera complanata	1
Caltha palustris	1	Oenanthe cf. aquatica	1
Campylium stellatum	1	oyster shell fgts	1 max 10 mm
Carex sp(p).	1 nutlets with utricles and/or free utricles	Polygonum aviculare agg.	1
		Polygonum lapathifolium	1
		Polygonum persicaria	1
cf. Avena sp(p). (chaff)	1	Polygonum persicaria/lapathifolium (ch)	1
cf. Calluna vulgaris (ch rt-tw fgts)	1	Prunella vulgaris	1
cf. Calluna vulgaris (rt-tw fgts)	1	Prunus domestica ssp. insititia	1
cf. Cladium mariscus (ch lf fgts)	1	Quercus (wood)	1 max 25mm
Chenopodium album (ch)	1	Quercus sp(p). (b/bs)	1
Chenopodium Section Pseudoblitum	1	Ranunculus flammula	1
Corylus avellana	1 including material with apical knife marks	Ranunculus sardous	1
		Ranunculus Section Ranunculus	1
		Raphanus raphanistrum (pod segs/fgts)	1
Corylus avellana (ch)	1	Rosa sp(p).	1
Cruciferae (pedicels)	1	Rubus fruticosus agg.	1
Daucus carota	1 very	Rumex sp(p).	1
decayed		Sambucus nigra	1
Diphasium complanatum	1	Sambucus nigra (ch)	1
Diphasium complanatum (ch)	1	sand	1
Drepanocladus sp(p).	1	Scorpidium scorpioides	1
earthworm egg caps	1	Sonchus asper	1
Eleocharis palustris sl	1	Stachys sp(p).	1
Euphorbia helioscopia	1	Stellaria media	1
Eurhynchium cf. striatum	1	stones	1 max 70mm
fish bone	1 max 10 mm	Thlaspi arvense	1
fly puparia	1	Thuidium cf. tamariscinum	1
fly puparia (min)	1	Triticum aestivo-compactum	1
Galeopsis Subgenus Galeopsis	1	unwashed peaty sediment	1
Galeopsis Subgenus Ladanum	1	Urtica urens	1
glassy slag	1 max 10 mm	Valerianella dentata	1
herbaceous detritus (ch)	1	Viola sp(p).	1
Homalothecium sericeum/lutescens	1	Viola sp(p). (caps segs)	1
Hordeum sp(p). (inc hulled)	1	woody root fgts (?modern)	1

Context 1005, Sample 10/T			
wood fgts	3 max 25mm	Legumino sae (pods/fgts)	1 max 5mm
Anthemis cotula	2	Leonto don sp(p).	1
bark fgts	2 max 40mm	Leucodon sciuroides	1
Chenopodium album	2	Linum usitatissimum	1 inc fgts
earthworm egg caps	2	Linum usitatissimum (caps fgts)	1
wood chips	2 max 5mm	Luzula sp(p).	1
Aethusa cynapium	1 fragment(s) only	Melophagus ovinus (sheep ked)	1
Agrostemma githago	1	Myrica gale (lf fgts)	1
Anthemis cotula (ch)	1	Picris hieracioides	1
Atriplex sp(p).	1	Polygonum aviculare agg.	1
Avena sativa (spklt/s/fgts)	1	Polygonum hydropiper	1
Avena sp(p).	1	Polygonum lapathifolium	1
Avena sp(p). (part-ch spklt/s/fgts)	1	Polygonum persicaria	1
beetles	1	Potentilla cf. erecta	1
Bilderdykia convolvulus	1	Potentilla palustris	1
bone fgts	1 max 15mm	pottery	1 max 15mm
burnt bone fgts	1 max 15mm	Prunella vulgaris	1
Calluna vulgaris (sht fgts)	1	Ranunculus Section Ranunculus	1
Campylium cf. elodes	1	Raphanus raphanistrum (pod segs/fgts)	1
Campylium stellatum	1	Rhytidiadelphus cf. squarrosus	1
Carex sp(p).	1	root/rootlet fgts	1
Centaurea sp(p).	1	Rubus fruticosus agg.	1
Cerealia indet.	1	Rumex acetosella agg.	1
cf. Avena sp(p). (chaff)	1	Rumex sp(p).	1
cf. Cladium mariscus (ch lf fgts)	1	Sambucus nigra	1
cf. Glyceria sp(p).	1	Satureja hortensis	1
charcoal	1 max 15mm	Scirpus setaceus	1
Cladium mariscus (with perig)	1	Scorpidium scorpioides	1
Corylus avellana	1	Secale cereale	1
dicot lf fgts	1	Stachys sp(p).	1
Diphasium complanatum	1	Stellaria media	1
Drepanocladus sp(p).	1	Thlaspi arvense	1
Eleocharis palustris sl	1	Thuidium tamariscinum	1
Eleocharis sp(p).	1	Triticum aestivo-compactum	1
Filipendula ulmaria	1	Urtica urens	1
fly puparia	1	Viola sp(p).	1
Fragaria cf. vesca	1	woody root fgts (?modern)	1
Fragaria cf. vesca (ch)	1		
Galeopsis Subgenus Ladanum	1	<hr/> Context 1011, Sample 7/SPT <hr/>	
Gramineae	1	Corylus avellana	1
Gramineae/Cerealia (c/n)	1		
Gramineae/Cerealia (ch c/n)	1	<hr/> Context 1013, Sample 16/SPT <hr/>	
herbaceous detritus	1	Corylus avellana	1
Hordeum sp(p).	1		
Humulus lupulus	1	<hr/> Context 1014, Sample 19/BS <hr/>	
Hylocomium cf. brevirostre	1	Chenopodium album	2
Hypnum cf. cupressiforme	1	Menyanthes trifoliata	2
Isatis tinctoria (pod fgts) decayed	1 very	Sambucus nigra	2
Isoetecium myurum	1	?cynipid galls	1
Juncus bufonius	1	Aethusa cynapium	1
Lamium Section Lamioipsis	1	Anthemis cotula	1
Lapsana communis	1	Atriplex sp(p).	1
leather fgts	1 max 5mm	Avena sp(p).	1
Legumino sae (fls/pet)	1		

beetles	1	Atriplex sp(p).	1
Bilderdykia convolvulus	1	Bilderdykia convolvulus	1
bone fgts	1 max 40mm	Bilderdykia convolvulus (ff)	1
Brassica rapa	1	bone fgts	1 max 20mm
Brassica sp./Sinapis arvensis	1	Brassica rapa	1
brick/tile	1 max 10 mm	Brassica sp./Sinapis arvensis	1
Bromus sp(p).	1	brick/tile	1 max 5mm
burnt bone fgts	1 max 10 mm	Bromus sp(p).	1
Cannabis sativa	1	Cannabis sativa	1 fragment(s) only
Carex sp(p).	1	Carduus/Cirsium sp(p).	1
cf. Secale cereale	1	cf. Avena sp(p).	1
charcoal	1 max 20mm	charcoal	1 max 5mm
Corylus avellana	1 including material with apical knife marks	Corylus avellana	1
Diphasium complanatum	1 very	Diphasium complanatum	1
decayed		earthworm egg caps	1
earthworm egg caps	1	Eleocharis palustris sl	1
Euphorbia helioscopia	1	fly puparia	1
Galeopsis Subgenus Galeopsis	1	Galeopsis Subgenus Galeopsis	1
gravel	1 max 25mm	Gramineae	1
Hordeum sp(p).	1	gravel	1 max 15mm
Humulus lupulus	1	Lapsana communis	1
Hyoscyamus niger	1	Linum usitatissimum	1 fragment(s) only
Lapsana communis	1	Melophagus ovinus (sheep ked)	1 fragment(s) only
Linum usitatissimum	1	Menyanthes trifoliata	1
mortar	1 max 5mm	Neckera complanata	1
Polygonum aviculare agg.	1	Nepeta cataria	1
Polygonum hydropiper	1	Polygonum persicaria	1
Polygonum lapathifolium	1	Ranunculus sardous	1
Prunus spinosa	1	Raphanus raphanistrum (pod segs/fgts)	1
Prunus spinosa (thorns)	1	root moulds (min)	1
Ranunculus sardous	1	Rubia tinctorum	1
Ranunculus Section Ranunculus	1	Rumex sp(p). (per/segs)	1
Rubus fruticosus agg.	1	Sambucus nigra	1
Rumex sp(p).	1	Scorpidium scorpioides	1
Scirpus maritimus	1	Sonchus asper	1
Scorpidium scorpioides	1	Sonchus oleraceus	1
Sonchus cf. arvensis	1	Stellaria media	1
textile fgts	1 max 5mm	Triglochin maritima	1
twig fgts	1	Triticum sp(p).	1 fragment(s) only
Urtica urens	1	twig fgts	1 max 20mm
Valerianella dentata	1	Urtica dioica	1
vivianite	1	Valerianella dentata	1
wood fgts	1 max 25mm	wood chips	1 max 5mm
		wood fgts	1 max 25mm

---

Context 1014, Sample 20/T

Anthemis cotula	2
bark fgts	2 max 45mm
Chenopodium album	2
concreted sediment	2 max 10 mm
Urtica urens	2
Aethusa cynapium	1
Agrostemma githago (sf)	1
Alnus glutinosa (b/bs)	1

---

Context 1030, Sample 22/BS

bark fgts	2 max 40mm
Aethusa cynapium	1
Agrostemma githago	1
Anagallis arvensis	1
Anthemis cotula	1
Atriplex sp(p).	1

Avena sp(p).	1	Sambucus nigra	1 inc fgts
beetles	1	Satureja hortensis	1
Betula cf. pendula (bark fgts)	1 max 25mm	Schoenus nigricans	1
Betula sp(p).	1	Scorpidium scorpioides	1
Betula sp(p). (fcs)	1	Sonchus asper	1
Bilderdykia convolvulus	1	Sphagnum sp(p). (lvs)	1 sp., not papillosum or imbricatum
bone fgts	1 max 15mm		
Brassica rapa	1		
Bromus sp(p).	1	Stellaria media	1
Calliargon cf. giganteum	1	Thuidium cf. tamariscinum	1
Cannabis sativa	1	Triticum aestivo-compactum	1
Carex sp(p).	1 nutlets with utricles and/or free utricles	Triticum sp(p).	1
		Triticum/Secale (w/l)	1 fragment(s) only
Centaurea sp(p).	1	twig fgts (min)	1 max 10 mm
cf. Cratoneuron filicinum	1	Urtica urens	1
cf. Rhytidiadelphus sp(p).	1	Valerianella dentata	1
cf. Secale cereale	1	Viola sp(p).	1
charcoal	1 max 25mm	wood fgts	1 max 30mm
Chenopodium album	1	wood fgts (min)	1 max 10 mm
Chenopodium murale	1		
cinders	1 max 10 mm		
coal	1 max 5mm	<hr/> Context 1030, Sample 24/T <hr/>	
Corylus avellana	1	Anthemis cotula	2
Danthonia decumbens	1	Chenopodium album	2
Diphasium complanatum	1	Dicranum scoparium	2
Drepanocladus sp(p).	1	grit	2
earthworm egg caps	1	Sambucus nigra	2 inc fgts
fly puparia	1	sand	2
Galeopsis Subgenus Galeopsis	1	Sphagnum sp(p). (lvs)	2 sp., not papillosum or imbricatum
Hordeum sp(p). (inc hulled)	1		
Isoethecium myurum	1		
Juglans regia	1		
Lamium Section Lamiopsis	1	?clumped sawdust	1 max 10 mm
Lapsana communis	1	?daub	1 max 25mm
Leguminosae (cal)	1	Aethusa cynapium	1
Leguminosae (fls/pet)	1	Alnus glutinosa	1
Leguminosae (pods/fgts)	1 max 5mm	Atriplex sp(p).	1
Leontodon sp(p).	1	Avena sp(p).	1
Linum usitatissimum	1 inc fgts	Avena sp(p). (chaff)	1
Malus sylvestris	1	Barbula sp(p).	1
Melophagus ovinus (sheep ked)	1	bark fgts	1 max 20mm
moss (contaminant)	1	beetles	1
Neckera complanata	1	Betula sp(p).	1
Pedicularis palustris	1	bone fgts	1 max 4mm
Pisum cf. sativum	1	Brassica rapa	1
Polygonum lapathifolium	1	Brassica sp(p).	1
Prunella vulgaris	1	brick/tile	1 max 40mm
Prunus spinosa (thorns)	1	burnt bone fgts	1 max 20mm
Quercus sp(p). (b/bs)	1	Carduus/Cirsium sp(p).	1
Ranunculus Section Ranunculus	1	Carex sp(p).	1
Raphanus raphanistrum (pod segs/fgts)	1	Ceratodon purpureus	1
Reseda luteola	1	cf. Amblystegium riparium	1
Rubus/Rosa sp(p). (prickles)	1	charcoal	1 max 10 mm
Rumex sp(p).	1	clinker	1 max 30mm
Rumex sp(p). (per/segs)	1	coal	1 max 30mm

<i>Corylus avellana</i>	1	bark fgts	1 max 25mm
Cruciferae	1 modern	<i>Betula</i> sp(p).	1 ?modern
<i>Diphysium complanatum</i>	1	<i>Betula</i> sp(p). (fcs)	1 ?modern
<i>Eleocharis palustris</i> sl (ch)	1	<i>Carex</i> sp(p).	1
<i>Erica tetralix</i> (lvs)	1	cf. <i>Calluna vulgaris</i> (ch rt-tw fgts)	1
<i>Ficus carica</i>	1	charcoal	1 max 10 mm
fish bone	1 max 4mm	<i>Chenopodium album</i>	1
fish scale	1 max 2mm	<i>Hordeum vulgare</i> (inc twstd)	1
fly puparia	1	<i>Linum usitatissimum</i> (ch)	1
Galeopsis Subgenus Galeopsis	1	<i>Polygonum aviculare</i> agg.	1
<i>Genista tinctoria</i> (st fgts)	1	<i>Raphanus raphanistrum</i> (ch pod segs/fgts)	1
Gramineae	1	root/rootlet fgts (modem)	1
gravel	1 max 15mm	<i>Triticum aestivo-compactum</i>	1
<i>Homalothecium sericeum/lutescens</i>	1	wood fgts	1 max 15mm
<i>Hylocomium splendens</i>	1		
<i>Hyoscyamus niger</i>	1		
<i>Hypnum</i> cf. <i>cupressiforme</i>	1	<hr/> Context 1034, Sample 31/T <hr/>	
<i>Hypochoeris</i> sp(p).	1	<i>Anthemis cotula</i>	3
<i>Isothecium myurum</i>	1	wood fgts	3 max 20mm
<i>Lamium</i> Section <i>Lamiopsis</i>	1	bark fgts	2 max 40mm
<i>Lapsana communis</i>	1	charcoal	2 max 10 mm
<i>Leucobryum glaucum</i>	1	<i>Chenopodium album</i>	2
<i>Leucodon sciuroides</i>	1	<i>Corylus avellana</i>	2
<i>Linum usitatissimum</i>	1 inc fgts	<i>Diphysium complanatum</i>	2 very
<i>Melophagus ovinus</i> (sheep ked)	1	decayed	
mortar	1 max 20mm	<i>Leontodon</i> sp(p).	2
<i>Neckera complanata</i>	1	<i>Valerianella dentata</i>	2
<i>Picris hieracioides</i>	1	'coils'	1
<i>Polygonum aviculare</i> agg.	1	?bees	1
<i>Populus</i> sp(p). (b/bs)	1	?daub	1 max 5mm
<i>Potentilla</i> cf. <i>erecta</i>	1	<i>Aethusa cynapium</i>	1
pottery	1 max 10 mm	<i>Agrostema githago</i> (sf)	1
<i>Prunus spinosa</i> (thorns)	1	<i>Anagallis arvensis</i>	1
<i>Pteridium aquilinum</i> (pinn fgts)	1	<i>Antitrichia curtispindula</i>	1
<i>Quercus</i> sp(p). (b/bs)	1	<i>Aphanes arvensis</i> agg.	1
<i>Ranunculus</i> Section <i>Ranunculus</i>	1	<i>Aster tripolium</i>	1
<i>Raphanus raphanistrum</i>	1	<i>Avena</i> sp(p).	1
<i>Raphanus raphanistrum</i> (pod segs/fgts)	1	<i>Avena</i> sp(p). (chaff)	1
<i>Rhytidiadelphus</i> sp(p).	1	bast fgts	1
root bark/epidermis fgts	1	bone fgts	1 max 5mm
<i>Rumex</i> sp(p).	1	<i>Brassica rapa</i>	1
<i>Schoenus nigricans</i>	1	<i>Brassica</i> sp(p).	1
<i>Scorpidium scorpioides</i>	1	<i>Brassica</i> sp./ <i>Sinapis arvensis</i>	1
<i>Silene alba</i>	1	brick/tile	1 max 4mm
small mammal tooth	1	burnt bone fgts	1 max 5mm
<i>Sonchus asper</i>	1	<i>Calliargon</i> cf. <i>giganteum</i>	1
<i>Sonchus palustris/arvensis</i>	1	<i>Carex</i> sp(p).	1
<i>Stellaria media</i>	1	<i>Centaurea</i> sp(p).	1
textile/yarn fgts	1	<i>Cladium mariscus</i> (with perig)	1
<i>Thuidium tamariscinum</i>	1	Compositae (inv fgts)	1
<i>Urtica urens</i>	1	concretions	1 max 10 mm
wood fgts	1 max 40mm	<i>Daucus carota</i>	1
		<i>Drepanoladus</i> sp(p).	1
		earthworm egg caps	1
<hr/> Context 1031, Sample 26/BS <hr/>		<i>Eleocharis palustris</i> sl	1
<i>Sambucus nigra</i>	2	<i>Eurhynchium striatum</i>	1
<i>Avena</i> sp(p).	1	fish bone	1 max 15mm



fly puparia	1	wood chips	2 max 10 mm
Galeopsis Subgenus Galeopsis	1	?amber	1 max 2mm
Gramineae	1	?daub	1 max 15mm
gravel	1 max 20mm	Aethusa cynapium	1
Homalothecium sericeum/lutescens	1	Atriplex sp(p).	1
Hordeum sp(p).	1	Atriplex sp(p). (ch)	1
Hyoscyamus niger	1	Avena sp(p).	1
Hypnum cf. cupressiforme	1	Avena sp(p). (part-ch spklt/sfgts)	1
Iris pseudacorus	1	beetles	1
Lapsana communis	1	Bilderdykia convolvulus	1
leather fgts	1 max 15mm	bone fgts	1 max 15mm
Leguminosae (fls/pet)	1	Brassica rapa	1 inc fgts
Leguminosae (imm s)	1	Brassica sp./Sinapis arvensis	1
Leguminosae (pods/fgts)	1	Brassica sp./Sinapis arvensis (min cot)	1
Leucodon sciuroides	1	brick/tile	1 max 10 mm
Linum usitatissimum	1 inc fgts	burnt bone fgts	1 max 15mm
Menyanthes trifoliata	1	Calliargon cuspidatum	1
Neckera complanata	1	Carex sp(p).	1
Polygonum aviculare agg.	1	Centaurea sp(p).	1
Polygonum hydropiper	1	Cerealia in det. (chaff)	1
Polygonum persicaria/lapathifolium	1	cf. Calluna vulgaris (ch rt-tw fgts)	1 max 5mm
Populus sp(p). (b/bs)	1	charred herbaceous detritus	1
Primulaceae	1	Corylus avellana	1
Prunella vulgaris	1	Cu/alloy fgts	1 very
Ranunculus Section Ranunculus	1	decayed	
Raphanus raphanistrum (pod segs/fgts)	1	Daucus carota	1
Rubia tinctorum	1 max 5mm	dicot lf fgts	1 max 2mm
Rubia tinctorum (root bark)	1	Diphysium complanatum	1 very
Rumex acetosella agg.	1	decayed	
sand	1	Eleocharis palustris sl	1
Satureja hortensis	1	fish bone	1 max 15mm
Schoenus nigricans	1	fish scale	1 max 2mm
sclereids (from bark)	1 max 1mm	fly puparia	1
Scorpidium scorpioides	1	gravel	1 max 15mm
sealing wax	1 max 5mm	Homalothecium sericeum/lutescens	1
Silene vulgaris	1	Hyoscyamus niger	1
Sonchus asper	1	Hypnum cf. cupressiforme	1
Sonchus palustris/arvensis	1	Hypochoeris sp(p).	1
Stellaria palustris/graminea	1	leather fgts	1 very
Thuidium tamariscinum	1		decayed,
Triticum/Secale (w/l)	1		max 5mm
twig fgts	1 max 20mm	Leontodon sp(p).	1
Viola sp(p).	1	Linum usitatissimum	1 inc fgts
wood chips	1 max 10 mm	Linum usitatissimum (caps fgts)	1
		Marrubium vulgare	1
		Menyanthes trifoliata	1
		mortar	1 max 5mm
		moss (contaminant)	1
		Neckera complanata	1
		Oxalis acetosella	1
		Plantago major	1
		Polygonum aviculare agg.	1
		Polygonum lapathifolium	1
		Polygonum persicaria	1
		pottery	1 max 35mm
		Prunella vulgaris	1
		Prunus spinosa (ch)	1 a single
<hr/>			
Context 103 5, Sample 32/T			
wood fgts	3 max 10 mm		
Anthemis cotula	2		
bark fgts	2 max 20mm		
charcoal	2 max 15mm		
Chenopodium album	2		
grit	2 max 1mm		
herbaceous detritus	2 mostly <1mm		
Quercus sp(p). (b/bs)	2		
sand	2		

	spec	dicot lf fgts	1
Ranunculus sardous	1	Diphasium complanatum	1 very
Ranunculus Section Ranunculus	1	decayed	
Raphanus raphanistrum (pod segs/fgts)	1	Drepanocladus sp(p).	1
Rhytidiadelphus sp(p).	1	earthworm egg caps	1
Rumex acetosella agg.	1	Eleocharis palustris sl	1
Rumex sp(p).	1	Eurhynchium striatum	1
Salix sp(p). (b)	1	fly puparia	1
Sambucus nigra	1 inc fgts	Genista tinctoria (st fgts)	1
sclereids (from bark)	1 max 2mm	Gramineae	1
Scorpidium scorpioides	1	Gramineae/Cerealia (c/n)	1
Secale cereale (rachis fgts)	1	gravel	1 max 25mm
Sonchus asper	1	grit	1
Sonchus oleraceus	1	Homalothecium sericeum/lutescens	1
Spergula arvensis	1	Hordeum sp(p).	1
Stellaria media	1	Hypnum cf. cupressiforme	1
Triticum aestivo-compactum	1	Isothecium myosuroides	1
twig fgts	1 max 10 mm	Lapsana communis	1
Ulota sp(p).	1	leather fgts	1 max 25mm
Urtica urens	1	Leguminosae (fls/pet)	1
Valerianella dentata	1	Leontodon sp(p).	1
Viola sp(p).	1	Leucodon sciuroides	1
		part-burnt wood	1 max 25mm
		Polygonum aviculare agg.	1
		Prunella vulgaris	1
		Prunus spinosa (thorns)	1
		Ranunculus flammula	1
		Ranunculus Section Ranunculus	1
		Raphanus raphanistrum (pod segs/fgts)	1
		Rhytidiadelphus sp(p).	1
		Rubia tinctorum	1 max 4mm
		sand	1
		sclereids (from bark)	1 max 2mm
		Sonchus asper	1
		Stellaria media	1
		Thuidium tamariscinum	1
		Triticum/Secale ('bran' fgts)	1 mostly <1mm
		twig fgts	1 max. 30 x 10 mm
		Ulota sp(p).	1
		undisaggreg. compressed plant debris	1 max 5mm
		Vicia faba (hila)	1
		wood chips	1 max 25mm
<hr/>			
Context 1037, Sample 29/SPT			
Corylus avellana	1 including		
material with apical knife marks			
<hr/>			
Context 1039, Sample 33/T			
bark fgts	3 max 40mm		
herbaceous detritus	3 mostly < 2mm		
wood fgts	3 max 45mm		
Anthemis cotula	2		
charcoal	2 max 25mm		
Juncus bufonius	2		
Linum usitatissimum	2 inc fgts		
Aethusa cynapium	1		
Agrostemma githago (sf)	1		
Alnus glutinosa (b/bs)	1		
Antitrichia curtipendula	1		
Aster tripolium	1		
Avena sp(p).	1		
Avena sp(p). (chaff)	1		
beetles	1		
Bilderdykia convolvulus (ff)	1		
bone fgts	1 max 80mm		
Brassica sp./Sinapis arvensis	1		
brick/tile	1 max 5mm		
Calliargon cf. giganteum	1		
Calliargon cuspidatum	1		
Carex sp(p).	1		
Chenopodium album	1		
concretions	1 max 10 mm		
Corylus avellana	1		

Table 4. Assemblages of adult Coleoptera and Hemiptera (excluding Aphidoidea and Coccidoidea) from the 41-49 Walmgate, York (Time Team) site: main statistics by sample. For explanation of codes see Table 6.

Context	1003	1004	1005	1014	1030	1034	1035	1039	Whole
Sample	11	12	10	20	24	31	32	33	site
Ext	/T	/T	/T	/T	/T	/T	/T	/T	
S	37	77	88	44	33	50	36	50	185
N	52	167	184	60	36	65	52	72	688
ALPHA	57	55	66	74	185	98	51	72	83
SEALPHA	16	7	8	20	107	28	14	17	5
SOB	9	17	17	14	6	16	8	13	65
PSOB	24	22	19	32	18	32	22	26	35
NOB	9	19	17	15	6	16	8	14	104
PNOB	17	11	9	25	17	25	15	19	15
ALPHAOB	0	0	0	0	0	0	0	0	74
SEALPHAOB	0	0	0	0	0	0	0	0	14
SW	1	2	2	1	0	2	1	3	7
PSW	3	3	2	2	0	4	3	6	4
NW	1	2	2	1	0	2	1	4	13
PNW	2	1	1	2	0	3	2	6	2
ALPHAW	0	0	0	0	0	0	0	0	0
SEALPHAW	0	0	0	0	0	0	0	0	0
SD	1	1	2	2	0	0	0	1	6
PSD	3	1	2	5	0	0	0	2	3
ND	1	1	2	2	0	0	0	1	7
PND	2	1	1	3	0	0	0	1	1
ALPHAD	0	0	0	0	0	0	0	0	0
SEALPHAD	0	0	0	0	0	0	0	0	0
SP	2	7	8	3	1	9	4	4	28
PSP	5	9	9	7	3	18	11	8	15
NP	2	8	8	4	1	9	4	4	40
PNP	4	5	4	7	3	14	8	6	6
ALPHAP	0	0	0	0	0	0	0	0	42
SEALPHAP	0	0	0	0	0	0	0	0	14
SM	0	0	0	0	0	0	0	0	0
PSM	0	0	0	0	0	0	0	0	0
NM	0	0	0	0	0	0	0	0	0
PNM	0	0	0	0	0	0	0	0	0
ALPHAM	0	0	0	0	0	0	0	0	0
SEALPHAM	0	0	0	0	0	0	0	0	0
SL	3	4	3	2	0	3	3	1	7
PSL	8	5	3	5	0	6	8	2	4
NL	3	4	3	2	0	3	3	1	19
PNL	6	2	2	3	0	5	6	1	3
ALPHAL	0	0	0	0	0	0	0	0	0

<b>Context</b>	<b>1003</b>	<b>1004</b>	<b>1005</b>	<b>1014</b>	<b>1030</b>	<b>1034</b>	<b>1035</b>	<b>1039</b>	<b>Whole</b>
<b>Sample</b>	<b>11</b>	<b>12</b>	<b>10</b>	<b>20</b>	<b>24</b>	<b>31</b>	<b>32</b>	<b>33</b>	<b>site</b>
<b>Ext</b>	<b>/T</b>	<b>/T</b>	<b>/T</b>	<b>/T</b>	<b>/T</b>	<b>/T</b>	<b>/T</b>	<b>/T</b>	
SEALPHAL	0	0	0	0	0	0	0	0	0
SRT	20	41	51	23	17	27	19	27	210
PSRT	54	53	58	52	52	54	53	54	114
NRT	35	117	135	37	19	40	35	41	459
PNRT	67	70	73	62	53	62	67	57	67
ALPHART	20	23	30	26	0	37	17	35	150
SEALPHART	6	3	4	8	0	12	5	11	12
SRD	5	11	12	4	5	7	6	6	56
PSRD	14	14	14	9	15	14	17	12	30
NRD	11	31	31	6	5	11	11	10	116
PNRD	21	19	17	10	14	17	21	14	17
ALPHARD	0	6	7	0	0	0	0	0	43
SEALPHARD	0	2	2	0	0	0	0	0	7
SRF	2	3	4	3	5	3	2	6	28
PSRF	5	4	5	7	15	6	6	12	15
NRF	3	7	4	3	6	5	3	10	41
PNRF	6	4	2	5	17	8	6	14	6
ALPHARF	0	0	0	0	0	0	0	0	39
SEALPHARF	0	0	0	0	0	0	0	0	13
SSA	16	33	42	17	11	19	13	22	62
PSSA	43	43	48	39	33	38	36	44	34
NSA	26	103	119	23	12	27	28	31	369
PNSA	50	62	65	38	33	42	54	43	54
ALPHASA	18	17	23	30	0	29	10	34	21
SEALPHASA	7	3	3	14	0	12	3	13	2
SSF	7	16	22	9	3	12	5	13	32
PSSF	19	21	25	20	9	24	14	26	17
NSF	9	31	40	12	4	15	6	17	134
PNSF	17	19	22	20	11	23	12	24	19
ALPHASF	0	14	20	0	0	0	0	0	13
SEALPHASF	0	4	6	0	0	0	0	0	2
SST	8	13	15	8	7	6	7	8	25
PSST	22	17	17	18	21	12	19	16	14
NST	16	66	59	11	7	11	13	13	196
PNST	31	40	32	18	19	17	25	18	28
ALPHAST	0	5	7	0	0	0	0	0	8
SEALPHAST	0	1	1	0	0	0	0	0	1
SSS	1	4	5	0	1	1	1	1	5
PSSS	3	5	6	0	3	2	3	2	3
NSS	1	6	20	0	1	1	9	1	39
PNSS	2	4	11	0	3	2	17	1	6
ALPHASS	0	0	2	0	0	0	0	0	2
SEALPHASS	0	0	1	0	0	0	0	0	0
SG	0	0	0	0	0	0	0	0	0

<b>Context</b>	<b>1003</b>	<b>1004</b>	<b>1005</b>	<b>1014</b>	<b>1030</b>	<b>1034</b>	<b>1035</b>	<b>1039</b>	<b>Whole</b>
<b>Sample</b>	<b>11</b>	<b>12</b>	<b>10</b>	<b>20</b>	<b>24</b>	<b>31</b>	<b>32</b>	<b>33</b>	<b>site</b>
<b>Ext</b>	<b>/T</b>	<b>/T</b>	<b>/T</b>	<b>/T</b>	<b>/T</b>	<b>/T</b>	<b>/T</b>	<b>/T</b>	
PSG	0	0	0	0	0	0	0	0	0
NG	0	0	0	0	0	0	0	0	0
PNG	0	0	0	0	0	0	0	0	0
ALPHAG	0	0	0	0	0	0	0	0	0
SEALPHAG	0	0	0	0	0	0	0	0	0

Table 5. Insects and other macro-invertebrates from 41-49 Walmgate, York (Time Team) site: species lists by sample. Taxa are listed in descending order of abundance then in taxonomic order. Key: n - minimum number of individuals; q - quantification (s - semi-quantitative 'several', m - semi-quantitative 'many', both sensu Kenward et al. (1986), e - estimate); ecodes - ecological codes (see Table 6); \* - not used in calculation of statistics in Table 4.

**Context: 1003 Sample: 11/T** ReM: S  
Weight: 2.00 E: 4.00 F: 3.00

Notes: Entered 28.2.2000. Recording approaching 'detail'. Flot 1 cm in jar, much 'char'. Possibly some unrecognisable charred invertebrates in addition to those listed. Recorded in flot, problems on filter paper. See sheet for details of charred remains.

Taxon	n	q	ecodes
Xylodromus concinnus	7	-	rt-st
Atomaria sp.	4	-	rd
Cryptophagus sp.	3	-	rd-sf
Platystethus arenarius	2	-	rf
Anotylus rugosus	2	-	rt
Cratarea suturalis	2	-	rt-st
Lathridius minutus group	2	-	rd-st
?Lyctocoris campestris	1	-	rd-st
Trechus quadristriatus	1	-	oa
Bembidion (Philochthus) sp.	1	-	oa
Bembidion sp.	1	-	oa
Carabidae sp.	1	-	ob
Helophorus sp.	1	-	oa-w
Cercyon analis	1	-	rt-sf
Platystethus cornutus group	1	-	oa-d
Anotylus nitidulus	1	-	rt
Leptacinus sp.	1	-	rt-st
Gyrophypnus angustatus	1	-	rt-st
Neobisnius sp.	1	-	u
Philonthus ?politus	1	-	rt-st
Staphylininae sp.	1	-	u
Falagria caesa or sulcatula	1	-	rt-sf
Aleocharinae sp. A	1	-	u
Aleocharinae sp. B	1	-	u
Aleocharinae sp. C	1	-	u
Euplectini sp.	1	-	u
Aphodius sp.	1	-	ob-rf
Anobium punctatum	1	-	l-sf
Ptinus ?fur	1	-	rd-sf
Omosita sp.	1	-	rt-sf
Corticaria sp.	1	-	rt-sf
Tenebrio obscurus	1	-	rt-ss
Salpingidae sp.	1	-	l
Anthicus floralis or formicarius	1	-	rt-st
Phymatodes alni	1	-	l
Longitarsus sp.	1	-	oa-p
Sitona sp.	1	-	oa-p
*Acarina sp.	15	m	u

*Oligochaeta sp. (egg capsule)	6	s	u
*Proctotrupoidea sp.	6	s	u
*Pulex irritans	2	-	ss
*Lepidoptera sp. (pupa)	1	-	u
*Diptera sp. (larva)	1	-	u
*Diptera sp. (puparium)	1	-	u
*Syrphidae sp. (larva)	1	-	u
*Coleoptera sp. (larva)	1	-	u
*Gastropoda sp.	1	-	u

**Context: 1004 Sample: 12/T** ReM: S  
Weight: 1.00 E: 3.50 F: 3.00

Notes: Entered 28/2/2000. Recording approaching 'detail'. Flot 6mm in jar. Recorded in flot and on filter paper. AH tube recorded: few remains. NB shrivelled Apion.

Taxon	n	q	ecodes
Xylodromus concinnus	29	-	rt-st
Lathridius minutus group	10	-	rd-st
Cercyon analis	8	-	rt-sf
Cratarea suturalis	8	-	rt-st
Aleocharinae sp. A	6	-	u
Cryptophagus scutellatus	6	-	rd-st
Aleocharinae sp. B	5	-	u
Platystethus arenarius	4	-	rf
Neobisnius sp.	4	-	u
Anotylus nitidulus	3	-	rt
Gyrophypnus angustatus	3	-	rt-st
Cordalia obscura	3	-	rt-sf
Ptinus ?fur	3	-	rd-sf
Atomaria nigripennis	3	-	rd-ss
Carpelimus ?bilineatus	2	-	rt-sf
Anotylus complanatus	2	-	rt-sf
Stenus sp. U	2	-	u
Gyrophypnus fracticornis	2	-	rt-st
Aphodius ?prodromus	2	-	ob-rf
Cryptophagus sp. A	2	-	rd-sf
Cryptophagus sp. B	2	-	rd-sf
Anthicus formicarius	2	-	rt-st
Phyllotreta nemorum group	2	-	oa-p
Lyctocoris campestris	1	-	rd-st
?Sphodrus leucophthalmus	1	-	oa
Bembidion sp.	1	-	oa
Pterostichus sp.	1	-	ob
Agonum dorsale	1	-	oa
Carabidae sp.	1	-	ob

Helophorus aquaticus or grandis	1	-	oa-w
Helophorus sp.	1	-	oa-w
Cercyon haemorrhoidalis	1	-	rf-sf
Acritus nigricornis	1	-	rt-st
Acrotichis sp.	1	-	rt
Catops sp.	1	-	u
Silpha atrata	1	-	u
Micropeplus sp.	1	-	rt
Phyllodrepa ?floralis	1	-	rt-sf
Dropephylla vilis	1	-	l
Omalium ?rivulare	1	-	rt-sf
Omalium sp.	1	-	rt
Coprophilus striatulus	1	-	rt-st
Carpelimus sp.	1	-	u
Anotylus rugosus	1	-	rt
Anotylus sculpturatus group	1	-	rt
Oxytelus sculptus	1	-	rt-st
Stenus sp. A	1	-	u
Leptacinus ?pusillus	1	-	rt-st
Philonthus ?politus	1	-	rt-st
Philonthus sp.	1	-	u
Quedius sp.	1	-	u
Staphylininae sp. A	1	-	u
Staphylininae sp. B	1	-	u
Tachinus laticollis or marginellus	1	-	u
Falagria caesa or sulcatula	1	-	rt-sf
Aleocharinae sp. C	1	-	u
Aleocharinae sp. D	1	-	u
Aleocharinae sp. E	1	-	u
Cyphon sp.	1	-	oa-d
Elateridae sp.	1	-	ob
Anobium punctatum	1	-	l-sf
Lyctus linearis	1	-	l-sf
Meligethes sp.	1	-	oa-p
Omosita ?discoidea	1	-	rt-sf
Atomaria sp. A	1	-	rd
Atomaria sp. B	1	-	rd
Ephistemus globulus	1	-	rd-sf
Corticaria sp.	1	-	rt-sf
Typhaea stercorea	1	-	rd-ss
Aglenus brunneus	1	-	rt-ss
Tenebrio obscurus	1	-	rt-ss
Cerambycidae sp.	1	-	l
Longitarsus sp.	1	-	oa-p
Chaetocnema ?concinna	1	-	oa-p
Apion (Oxystoma) ?subulatum	1	-	oa-p
Apion sp.	1	-	oa-p
Ceutorhynchus sp.	1	-	oa-p
*Proctotrupoidea sp.	15	m	u
*Acarina sp.	15	m	u
*Pulex irritans	8	-	ss
*Oligochaeta sp. (egg capsule)	6	s	u
*Diptera sp. (puparium)	6	s	u
*Coleoptera sp. (larva)	6	s	u
*Aranae sp.	6	-	u
*Auchenorhyncha sp. (nymph)	3	-	oa-p

*Melophagus ovinus (adult)	2	-	oa-w
*Melophagus ovinus (puparium)	1	-	u
*Syrphidae sp. (larva)	1	-	u
*Chalcidoidea sp.	1	-	u
*Insecta sp. (larva)	1	-	u

**Context: 1005 Sample: 10/T ReM: S**  
Weight: 1.00 E: 2.50 F: 2.50

Notes: Entered 28.2.2000. Recording approaching 'detail'. Flot 1 cm in jar, remains part sorted to filter paper during Time Team extravaganza. Recorded in flot and on filter paper. Aleocharinae represent a minimum of the taxa present

Taxon	n	q	ecodes
Xylodromus concinnus	20	-	rt-st
Aglenus brunneus	14	-	rt-ss
Cryptophagus scutellatus	8	-	rd-st
Cercyon analis	6	-	rt-sf
Ptenidium sp.	5	-	rt
Leptacinus pusillus	5	-	rt-st
Aleocharinae sp. A	5	-	u
Gyrohypnus fracticornis	4	-	rt-st
Cryptophagus sp. B	4	-	rd-sf
Lathridius minutus group	4	-	rd-st
Micropeplus fulvus	3	-	rt
Omalium caesum or italicum	3	-	rt-sf
Carpelimus fuliginosus	3	-	st
Oxytelus sculptus	3	-	rt-st
Cratarea suturalis	3	-	rt-st
Atomaria nigripennis	3	-	rd-ss
Lyctocoris campestris	2	-	rd-st
Catops sp.	2	-	u
Omalium excavatum	2	-	rt-sf
Carpelimus bilineatus	2	-	rt-sf
Anotylus nitidulus	2	-	rt
Stenus sp.	2	-	u
Neobisnius sp.	2	-	u
Cordalia obscura	2	-	rt-sf
Aleocharinae sp. B	2	-	u
Aleocharinae sp. E	2	-	u
Pselaphidae sp.	2	-	u
Clambus ?pubescens	2	-	rt-sf
Ptinus ?fur	2	-	rd-sf
Monotoma spinicollis	2	-	rt-st
Cryptophagus sp. A	2	-	rd-sf
Atomaria sp. A	2	-	rd
Orthoperus sp.	2	-	rt
Corticaria sp. B	2	-	rt-sf
Empicoris culiciformis	1	-	rt
Chartoscirta ?cocksii	1	-	oa-w
Auchenorhyncha sp.	1	-	oa-p
Bembidion sp.	1	-	oa
Pterostichus (Poecilus) sp.	1	-	oa
Odacantha melanura	1	-	oa-d

Carabidae sp.	1	-	ob
Cercyon haemorrhoidalis	1	-	rf-sf
Cercyon unipunctatus	1	-	rf-st
Acritus nigricornis	1	-	rt-st
Ochthebius ?minus	1	-	oa-w
Acrotichis sp.	1	-	rt
?Phyllodrepa salicis	1	-	rt-sf
Platystethus arenarius	1	-	rf
Platystethus degener	1	-	oa-d
Anotylus complanatus	1	-	rt-sf
Anotylus rugosus	1	-	rt
Phacophallus parumpunctatus	1	-	rt-st
Philonthus ?politus	1	-	rt-st
Philonthus sp. A	1	-	u
Philonthus sp. B	1	-	u
Creophilus maxillosus	1	-	rt
Quedius sp.	1	-	u
Staphylininae sp. A	1	-	u
Staphylininae sp. B	1	-	u
Tachinus laticollis or marginellus	1	-	u
Falagria caesa or sulcatula	1	-	rt-sf
Aleocharinae sp. C	1	-	u
Aleocharinae sp. D	1	-	u
Aleocharinae sp. F	1	-	u
Aleocharinae sp. G	1	-	u
Aphodius sp.	1	-	ob-rf
Clambus sp.	1	-	rt-sf
Elateridae sp.	1	-	ob
Anobium punctatum	1	-	l-sf
Lyctus linearis	1	-	l-sf
Meligethes sp.	1	-	oa-p
Monotoma sp.	1	-	rt-sf
Cryptophagus sp. C	1	-	rd-sf
Atomaria sp. C	1	-	rd
Mycetaea hirta	1	-	rd-ss
Enicmus sp.	1	-	rt-sf
Corticaria sp. A	1	-	rt-sf
Typhaea stercorea	1	-	rd-ss
Tenebrio obscurus	1	-	rt-ss
Salpingidae sp.	1	-	l
Anthicus floralis or formicarius	1	-	rt-st
Phaedon sp.	1	-	oa-p
Phyllotreta sp.	1	-	oa-p
Chaetocnema sp.	1	-	oa-p
Apion sp.	1	-	oa-p
Dorytomus sp.	1	-	oa-p
Ceutorhynchus contractus	1	-	oa-p
*Acarina sp.	100	e	u
*Oligochaeta sp. (egg capsule)	15	m	u
*Proctotrupoidea sp.	15	m	u
*Diptera sp. (pupa)	6	s	u
*Diptera sp. (puparium)	6	s	u
*Araneae sp.	6	-	u
*Pulex irritans	3	-	ss
*Coccoidea sp.	2	-	u
*Hymenoptera sp.	2	-	u

*Daphnia sp. (ephippium)	1	-	oa-w
*Cladocera sp. (ephippium)	1	-	oa-w
*Heteroptera sp. (nymph)	1	-	u
*Auchenorhyncha sp. (nymph)	1	-	oa-p
*Coleoptera sp. (larva)	1	-	u

**Context: 1014 Sample: 20/T** ReM: S

Weight: 1.00 E: 3.00 F: 3.00

Notes: Entered 28/2/2000. Recording approaching 'detail'. Flot 6-8mm in jar, very coarse. Recorded in flot, problems on filter paper. AH tube listed: little of consequence.

Taxon	n	q	ecodes
Anotylus nitidulus	6	-	rt
Cercyon analis	3	-	rt-sf
Xylodromus concinnus	3	-	rt-st
Anotylus rugosus	3	-	rt
Trechus micros	2	-	u
Gyrophynus fracticornis	2	-	rt-st
Cryptophagus sp.	2	-	rd-sf
Atomaria sp.	2	-	rd
Ceutorhynchus contractus	2	-	oa-p
Clivina fossor	1	-	oa
Bembidion sp.	1	-	oa
Stomis pumicatus	1	-	oa
Carabidae sp.	1	-	ob
Hydraena sp.	1	-	oa-w
Omalium caesum or italicum	1	-	rt-sf
Omalium ?rivulare	1	-	rt-sf
Platystethus ?degener	1	-	oa-d
Platystethus nitens	1	-	oa-d
Anotylus complanatus	1	-	rt-sf
Oxytelus sculptus	1	-	rt-st
Stenus sp.	1	-	u
Gyrophynus angustatus	1	-	rt-st
Neobisnius sp.	1	-	u
Falagria caesa or sulcatula	1	-	rt-sf
Cratarea suturalis	1	-	rt-st
Aleocharinae sp. A	1	-	u
Aleocharinae sp. B	1	-	u
Aleocharinae sp. C	1	-	u
Staphylinidae sp.	1	-	u
Euplectini sp.	1	-	u
Aphodius granarius	1	-	ob-rf
Aphodius prodromus	1	-	ob-rf
Aphodius sp.	1	-	ob-rf
Hoplia philanthus	1	-	oa
Anobium punctatum	1	-	l-sf
Omosita sp.	1	-	rt-sf
Cryptophagus scutellatus	1	-	rd-st
Orthoperus sp.	1	-	rt
Lathridius minutus group	1	-	rd-st
Corticaria sp.	1	-	rt-sf
Anthicus floralis or formicarius	1	-	rt-st



Cerambycidae sp.	1	-	l
Longitarsus sp.	1	-	oa-p
Apion sp.	1	-	oa-p
*Acarina sp.	15	m	u
*Oligochaeta sp. (egg capsule)	6	s	u
*Proctotrupoidea sp.	3	-	u
*Diptera sp. (puparium)	2	-	u
*Pulex irritans	2	-	ss
*Coleoptera sp. (larva)	2	-	u
*Aranae sp.	2	-	u
*?Trichoptera sp.	1	-	oa-w
*Melophagus ovinus (puparium)	1	-	u
*Hymenoptera Parasitica sp.	1	-	u
*Insecta sp. (larva)	1	-	u

**Context: 1030 Sample: 24/T** ReM: S  
Weight: 1.00 E: 3.50 F: 3.50

Notes: Entered 28/2/2000. One dish flot. Recorded in flot, problems on filter paper.

Taxon	n	q	ecodes
Carpelimus ?bilineatus	2	-	rt-sf
Carpelimus pusillus group	2	-	u
Platystethus arenarius	2	-	rf
?Hemiptera sp.	1	-	u
?Trechus sp.	1	-	ob
Carabidae sp.	1	-	ob
Cercyon analis	1	-	rt-sf
Cercyon unipunctatus	1	-	rf-st
Cryptopleurum minutum	1	-	rf-st
Acritus nigricornis	1	-	rt-st
Carpelimus fuliginosus	1	-	st
Anotylus nitidulus	1	-	rt
Anotylus rugosus	1	-	rt
Oxytelus sculptus	1	-	rt-st
Stenus sp. A	1	-	u
Stenus sp. B	1	-	u
Lathrobium sp.	1	-	u
Neobisnius sp.	1	-	u
Philon thus sp. A	1	-	u
Philon thus sp. B	1	-	u
Falagria or Cordalia sp.	1	-	rt-sf
Aleocharinae sp. A	1	-	u
Aleocharinae sp. B	1	-	u
Aphodius ?granarius	1	-	ob-rf
Aphodius sp.	1	-	ob-rf
Ptinidae sp.	1	-	rd
Rhizophagus sp.	1	-	u
Cryptophagus scutellatus	1	-	rd-st
Atomaria sp.	1	-	rd
Mycetaea hirta	1	-	rd-ss
Lathridius minutus group	1	-	rd-st
Chaetocnema sp.	1	-	oa-p
Curculionidae sp.	1	-	oa

*Acarina sp.	15	m	u
*Proctotrupoidea sp.	3	-	u
*Diptera sp. (puparium)	2	-	u
*Cladocera sp. F (ephippium)	1	-	oa-w
*Melophagus ovinus (puparium)	1	-	u
*Siphonaptera sp.	1	-	u
*Coleoptera sp. (larva)	1	-	u
*Apoidea sp.	1	-	u
*Formicidae sp.	1	-	u

**Context: 1034 Sample: 31/T** ReM: S  
Weight: 1.00 E: 3.50 F: 3.50

Notes: Entered 28/2/2000. Flot 5mm in jar. Recorded in flot, problems on filter paper. AH tube listed: inconsequential. Subjectively, hints of heath/mo or.

Taxon	n	q	ecodes
Xylodromus concinnus	4	-	rt-st
Carpelimus pusillus group	3	-	u
Platystethus arenarius	3	-	rf
Omalium caesum or italicum	2	-	rt-sf
Anotylus complanatus	2	-	rt-sf
Anotylus nitidulus	2	-	rt
Anotylus rugosus	2	-	rt
Cryptophagus scutellatus	2	-	rd-st
Cryptophagus sp. B	2	-	rd-sf
Atomaria sp. B	2	-	rd
Lathridius minutus group	2	-	rd-st
Delphacidae sp. A	1	-	oa-p
Delphacidae sp. B	1	-	oa-p
Auchenorhyncha sp.	1	-	oa-p
Amara sp.	1	-	oa
Carabidae sp.	1	-	ob
Helophorus sp.	1	-	oa-w
Cercyon analis	1	-	rt-sf
Histerinae sp.	1	-	rt
Ochthebius sp.	1	-	oa-w
Acrotichis sp.	1	-	rt
Phyllodrepa ?floralis	1	-	rt-sf
Dropephylla sp.	1	-	u
Omalium ?rivulare	1	-	rt-sf
Carpelimus ?bilineatus	1	-	rt-sf
Stenus sp.	1	-	u
Gyrophypnus angustatus	1	-	rt-st
Gyrophypnus fracticornis	1	-	rt-st
Falagria or Cordalia sp.	1	-	rt-sf
Aleocharinae sp. A	1	-	u
Aleocharinae sp. B	1	-	u
Aleocharinae sp. C	1	-	u
Trox scaber	1	-	rt-sf
Aphodius sp. A	1	-	ob-rf
Aphodius sp. B	1	-	ob-rf
Elateridae sp.	1	-	ob
Anobium punctatum	1	-	l-sf
Ptinus sp.	1	-	rd-sf

Cryptophagus sp. A	1	-	rd-sf	Anotylus nitidulus	1	-	rt
Atomaria sp. A	1	-	rd	Stenus sp.	1	-	u
Aglenus brunneus	1	-	rt-ss	Gyrophypnus angustatus	1	-	rt-st
Anthicus floralis or formicarius	1	-	rt-st	Gyrophypnus ?fracticornis	1	-	rt-st
Phymatodes alni	1	-	l	Philonthus sp. A	1	-	u
Cerambycidae sp.	1	-	l	Philonthus sp. B	1	-	u
?Altica sp.	1	-	oa-p	Crataraea suturalis	1	-	rt-st
Halticinae sp.	1	-	oa-p	Aleocharinae sp. A	1	-	u
Apion sp. A	1	-	oa-p	Aleocharinae sp. B	1	-	u
Apion sp. B	1	-	oa-p	Trox scaber	1	-	rt-sf
?Strophosomus sp.	1	-	oa-p	Aphodius sp.	1	-	ob-rf
?Sitona sp.	1	-	oa-p	Lyctus linearis	1	-	l-sf
				Atomaria sp. A	1	-	rd
*Acarina sp.	15	m	u	Atomaria sp. B	1	-	rd
*Diptera sp. (puparium)	6	s	u	Orthoperus sp.	1	-	rt
*Proctotrupoidea sp.	6	s	u	Cerambycidae sp.	1	-	l
*Pulex irritans	4	-	ss	Longitarsus sp.	1	-	oa-p
*Oligochaeta sp. (egg capsule)	3	-	u	Chaetocnema sp.	1	-	oa-p
*Coccoidea sp.	3	-	u	Cossoninae sp.	1	-	u
*Diptera sp. (adult)	2	-	u	Ceuthorhynchinae sp.	1	-	oa-p
*Forficula auricularia	1	-	rt	Leperisinus ?varius	1	-	l
*Auchenomyncha sp. (nymph)	1	-	oa-p				
*Diptera sp. (pupa)	1	-	u	*Acarina sp.	15	m	u
*Melophagus ovinus (adult)	1	-	oa-w	*Coccoidea sp.	3	-	u
*Melophagus ovinus (puparium)	1	-	u	*Oligochaeta sp. (egg capsule)	2	-	u
*Syrphidae sp. (larva)	1	-	u	*Auchenomyncha sp. (nymph)	2	-	oa-p
*Dermestes lardarius (larva)	1	-	rd-sf	*Cladocera sp. (ephippium)	1	-	oa-w
*?Apis mellifera	1	-	u	*Syrphidae sp. (larva)	1	-	u
*Chalcidoidea sp.	1	-	u	*Siphonaptera sp.	1	-	u
*Gastropoda sp.	1	-	u	*Insecta sp. (larva)	1	-	u
*Lophopus crystallinus	1	-	oa-w				

**Context: 1035 Sample: 32/T** ReM: S  
Weight: 1.00 E: 3.50 F: 2.50

Notes: Entered 29.2.2000. Two dish flot. Recorded in flot, problems on filter paper. Recording approaching 'detail'. AH tube, with few useful remains. Entire modern Carpelimus gracilis to reference collection.

Taxon	n	q	ecodes	Taxon	n	q	ecodes
Aglenus brunneus	9	-	rt-ss	Platystethus arenarius	5	-	rf
Cryptophagus scutellatus	4	-	rd-st	Carpelimus pusillus group	4	-	u
Xylodromus concinnus	3	-	rt-st	Anotylus complanatus	4	-	rt-sf
Platystethus arenarius	2	-	rf	Lathridius minutus group	4	-	rd-st
Cryptophagus sp.	2	-	rd-sf	Stenus sp. U	3	-	u
Lathridius minutus group	2	-	rd-st	Aleocharinae sp. B	3	-	u
?Lyctocoris campestris	1	-	rd-st	Helophorus sp.	2	-	oa-w
?Paramesus nervosus	1	-	oa-p	Xylodromus concinnus	2	-	rt-st
Carabidae sp. A	1	-	ob	Anotylus nitidulus	2	-	rt
Carabidae sp. B	1	-	ob	Gyrophypnus ?fracticornis	2	-	rt-st
Helophorus ?aquaticus	1	-	oa-w	Cryptophagus sp. A	2	-	rd-sf
Cercyon analis	1	-	rt-sf	Aphrodes sp.	1	-	oa-p
Ptenidium sp.	1	-	rt	Delphacidae sp.	1	-	oa-p
Carpelimus ?bilineatus	1	-	rt-sf	Bembidion sp.	1	-	oa
Carpelimus pusillus group	1	-	u	Helophorus aquaticus or grandis	1	-	oa-w
				Cercyon analis	1	-	rt-sf
				Cercyon ?unipunctatus	1	-	rf-st

**Context: 1039 Sample: 33/T** ReM: S  
Weight: 1.00 E: 3.50 F: 2.50

Notes: Entered 28/2/2000. Recording approaching 'detail'. Small flot. Recorded in flot, problems on filter paper. AH tube contained no significant remains.

Hydraena sp.	1	-	oa-w
Scydmaenus sp.	1	-	u
Dropephylla ?ioptera	1	-	u
Omalium caesum or italicum	1	-	rt-sf
Omalium ?rivulare	1	-	rt-sf
Coprophilus striatulus	1	-	rt-st
Carpelimus bilineatus	1	-	rt-sf
Carpelimus fuliginosus	1	-	st
Platystethus nitens	1	-	oa-d
Leptacinus pusillus	1	-	rt-st
Philonthus sp.	1	-	u
Staphylininae sp.	1	-	u
Falagria or Cordalia sp.	1	-	rt-sf
Aleocharinae sp. A	1	-	u
Aleocharinae sp. C	1	-	u
Aleocharinae sp. D	1	-	u
Aleocharinae sp. E	1	-	u
Euplectini sp.	1	-	u
Aphodius granarius	1	-	ob-rf
Aphodius ?prodromus	1	-	ob-rf
Aphodius sp. A	1	-	ob-rf
Aphodius sp. B	1	-	ob-rf
Anobium punctatum	1	-	l-sf
Ptinus sp.	1	-	rd-sf
?Meligethes sp.	1	-	oa-p
Omosita sp.	1	-	rt-sf
Cryptophagus ?scutellatus	1	-	rd-st
Cryptophagus sp. B	1	-	rd-sf
Atomaria sp.	1	-	rd
Corticaria sp. A	1	-	rt-sf
Corticaria sp. B	1	-	rt-sf
Aglenus brunneus	1	-	rt-ss
?Longitarsus sp.	1	-	oa-p
*Diptera sp. (puparium)	300	e	u
*Acarina sp.	15	m	u
*Diptera sp. (pupa)	6	s	u
*Auchenorhyncha sp. (nymph)	3	-	oa-p
*Coccoidea sp.	3	-	u
*Coleoptera sp. (larva)	3	-	u
*Diptera sp. (adult)	2	-	u
*Oligochaeta sp. (egg capsule)	1	-	u
*Daphnia sp. (ephippium)	1	-	oa-w
*Melophagus ovinus (adult)	1	-	oa-w
*Siphonaptera sp.	1	-	u
*Chalcidoidea sp.	1	-	u
*Gastropoda sp.	1	-	u

Table 6. Abbreviations for ecological codes and statistics used for interpretation of insect remains in text and tables. Lower case codes in parentheses are those assigned to taxa and used to calculate the group values (the codes in capitals). See Table 2 for codes assigned to taxa from 41-49 Walmgate, York. Alpha - the index of diversity alpha (Fisher et al. 1943); Indivs - individuals (based on MNI); No - number.

No taxa	S	Percentage of indivs of grain pests	PNG
Estimated number of indivs (MNI)	N	No decomposer taxa (rt + rd + rf)	SRT
Index of diversity ( )	alpha	Percentage of RT taxa	PSRT
Standard error of alpha	SE alpha	No RT indivs	NRT
No 'certain' outdoor taxa (oa)	SOA	Percentage of RT indivs	PNRT
Percentage of 'certain' outdoor taxa	PSOA	Index of diversity of RT component	alpha RT
No 'certain' outdoor indivs	NOA	Standard error	SEalphaRT
Percentage of 'certain' outdoor indivs	PNOA	No 'dry' decomposer taxa (rd)	SRD Percentage
No OA and probable outdoor taxa (oa+ob)	SOB	of RD taxa	PSRD
Percentage of OB taxa	PSOB	No RD indivs	NRD
No OB indivs	NOB	Percentage of RD indivs	PNRD
Percentage OB indivs	PNOB	Index of diversity of the RD component	alphaRD
Index of diversity of the OB component	alphaOB	Standard error	SEalphaRD
Standard error	SEalphaOB	No 'foul' decomposer taxa (rf)	SRF
No aquatic taxa (w)	SW	Percentage of RF taxa	PSRF
Percentage of aquatic taxa	PSW	No RF indivs	NRF
No aquatic indivs	NW	Percentage of RF indivs	PNRF
Percentage of W indivs	PNW	Index of diversity of the RF component	alphaRF
Index of diversity of the W component	alphaW	Standard error	SEalphaRF
Standard error	SEalphaW	No synanthropic taxa (sf+st_ss)	SSA
No damp ground/waterside taxa (d)	SD	Percentage of synanthropic taxa	PSSA
Percentage D taxa	PSD	No synanthropic indivs	NSA
No damp D indivs	ND	Percentage of SA indivs	PNSA
Percentage of D indivs	PND	Index of diversity of SA component	ALPHASA
Index of diversity of the D component	alphaD	Standard error	SEALPHASA
Standard error	SEalphaD	No facultatively synanthropic indivs	SSF
No strongly plant-associated taxa (p)	SP	Percentage of SF taxa	PSSF
Percentage of P taxa	PSP	No SF indivs	NSF
No strongly P indivs	NP	Percentage of SF indivs	PNSF
Percentage of P indivs	PNP	Index of diversity of SF component	ALPHASF
Index of diversity of the P component	alphaP	Standard error	SEALPHASF
Standard error	SEalphaP	No typical synanthropic indivs	SST
No heathland/moorland taxa (m)	SM	Percentage of ST taxa	PSST
Percentage of M taxa	PSM	No ST indivs	NST
No M indivs	NM	Percentage of ST indivs	PNST
Percentage of M indivs	PNM	Index of diversity of ST component	ALPHAST
Index of diversity of the M component	alphaM	Standard error	SEALPHAST
Standard error	SEalphaM	No strongly synanthropic taxa	SSS
No wood-associated taxa (l)	SL	Percentage of SS taxa	PSSS
Percentage of L taxa	PSL	No SS indivs	NSS
No L indivs	NL	Percentage of SS indivs	PNSS
Percentage of L indivs	PNL	Index of diversity of SS component	ALPHASS
Index of diversity of the L component	alphaL	Standard error	SEALPHASS
Standard error	SEalphaL	No uncoded taxa (u)	SU
No indivs of grain pests (g)	NG	Percentage of uncoded indivs	PNU

Table 7. 41-49 Walmgate, York (Time Team) site. Preservation records for assemblages of macro-invertebrates. Key: E - degree of chemical erosion; F - degree of fragmentation (in each case 0.5 - none ... 5.5 - extreme). For strength (of mode): W - weak; D - distinct; S - strong. See Kenward and Large (1998) for full explanation.

Context	Sample	E			F			Other
		Range	Mode	Strength	Range	Mode	Strength	
1003	11	3.0-5.0	4.0	W	2.0-5.0	3.0	W	Numerous charred remains present
1004	12	1.5-4.0	3.5	D	2.0-5.0	3.0	D	Trace of charred remains
1005	10	1.5-3.5	2.5	W	1.5-4.0	2.5	D	Crumpled <i>Pulex</i> head
1014	20	1.5-4.0	3.0	D	1.0-4.0	3.0	D	
1030	24	3.0-4.0	3.5	W	2.5-5.5	3.5	W	
1034	31	3.0-4.5	3.5	W	2.0-5.0	3.5	W	
1035	32	2.0-4.0	3.5	D	1.5-4.5	2.5	D	
1039	33	2.0-4.5	3.5	D	1.5-4.5	2.5	S	

Table 8. Assemblages of adult Coleoptera and Hemiptera (excluding Aphidoidea and Coccidoidea) from 41-49 Walmgate, York (Time Team) site: Numbers of taxa (s) and individuals (n) placed in Group A ('house fauna') by Carrott and Kenward (2000), by sample. Note that the various fleas, lice and flies placed in Group A are excluded. The 'core' group consists of taxa with strong affinities only within the group: 'shared' taxa have some affinities with members of other groups.

Context	Sample	Ext	N	s (core)	n (core)	n (core) as % of N	s (core + shared)	n (core + shared)	n (core + shared) as % of N
1003	11	/T	52	10	23	44	12	25	48
1004	12	/T	167	20	75	45	22	79	47
1005	10	/T	184	22	77	42	29	89	48
1014	20	/T	60	8	12	20	10	14	23
1030	24	/T	36	4	4	11	5	5	14
1034	31	/T	65	10	17	26	12	20	31
1035	32	/T	52	11	26	50	11	26	50
1039	33	/T	72	10	15	21	13	18	25

Table 9. 41-49 Walmgate, York (Time Team) site. Charred invertebrate remains from Sample 11/T, Context 1003. Key: *abd* - abdomen; *hd* - head capsule; *th* - prothorax or pronotum; *us* - thoracic sterna.

Taxon	Charred parts recorded
Diptera puparia	fragment
<i>Pulex irritans</i>	entire including most of appendages
Carabidae sp.	leg
<i>Cercyon analis</i>	?us+abd
<i>Xylodromus concinnus</i>	one entire
	hd (with antennae)+th+us
	hd
Euplectinae sp.	entire apart from appendages
<i>Cryptophagus</i> sp.	?us+abd
<i>Atomaria</i> sp.	hd+th+abd
	us+abd
Proctotrupoidea sp.	hd
Acarina	several entire apart from distal parts of limbs

Table 10. Summary of the shell remains recovered from well-dated contexts at 41-49 Walmgate, York (numbers of hand-collected and SRS recovered combined by Context). A '?' before numbers indicates possible numbers (e.g. '3(?4)' = definitely 3, possibly 4, whereas '?3/4' = possibly 3, possibly 4).

**Key (Oyster):** 'Rv' = number of right (or upper) valves; 'Lv' = number of left (or lower) valves; 'Iv' = number of valves of indeterminate side; 'Knife marks' = number of valves showing damage characteristic of the oyster having been opened using a knife or similar implement; '?Meas' = estimated number of valves intact enough to be measured usefully; 'Worm' = number of valves showing damage by polychaet worms; 'Barn' = number of valves with barnacles; 'Dog whelk' = number of valves showing damage from dog whelk boring.

Context	<i>Helix</i> sp.	Mussel	Oyster							
			Rv	Lv	Iv	Knife marks	?Meas	Worm	Barn	Dog whelk
1013	1	0	5	8	0	1(?3)	3(?4)	0	0	0
1030	0	1	2	3	0	1	4	0	0	0
1031	0	0	0	0	?1	0	0	0	0	0
<b>Total</b>	<b>1</b>	<b>1</b>	<b>7</b>	<b>11</b>	<b>?1</b>	<b>2(?4)</b>	<b>7(?8)</b>	<b>0</b>	<b>0</b>	<b>0</b>

Table 11. Total numbers of vertebrate fragments, with numbers of measurable and subadult bones, mandibles and isolated teeth yielding ageing and sexing information and weights, by species, from 41-49 Walmgate (Time Team site), York.

Key: Total frags = total number of fragments; No. meas = number of measurable fragments; No. mand = number of mandibles; No. teeth = number of isolated mandibular teeth; No. unfused = number of unfused fragments.

Taxon		No. meas	No. unfused	No. mand	No. teeth	Total frags	Weight (g)
Vole/ mouse	Microtine/Murine	-	-	-	-	1	0.1
Bank vole	<i>Clethrionomys glareolus</i> (Schreber)	-	-	-	-	1	0.1
Mouse	Murine sp.	-	-	-	-	1	0.1
Cat	<i>Felis f. domestic</i>	1	-	-	-	1	1.3
Pig	<i>Sus f. domestic</i>	-	-	1	1	16	152.2
Cattle	<i>Bos f. domestic</i>	1	1	1	1	24	636.8
Sheep/ goat	Caprovid	3	1	-	1	12	64.5
Duck	<i>Anas sp.</i>	1	-	-	-	1	1.7
?Fowl	cf. <i>Gallus f. domestic</i>	1	-	-	-	1	0.5
Chicken	<i>Gallus f. domestic</i>	2	-	-	-	2	7.2
Herring	<i>Clupea harengus L.</i>	-	-	-	-	25	0.6
Eel	<i>Anguilla anguilla</i> (L.)	-	-	-	-	5	0.4
Cod family	Gadid	-	-	-	-	3	2.9
Cod	<i>Gadus morhua L.</i>	-	-	-	-	9	9.7
Haddock	<i>Melanogrammus aeglefinus</i> (L.)	-	-	-	-	2	0.7
<b>Subtotal</b>		<b>9</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>104</b>	<b>878.8</b>
Large mammal		-	-	-	-	126	1377.4
Medium-sized mammal 1		-	-	-	-	164	377.7

<b>Taxon</b>		<b>No. meas</b>	<b>No. unfused</b>	<b>No. mand</b>	<b>No. teeth</b>	<b>Total frags</b>	<b>Weight (g)</b>
Bird		-	-	-	-	20	7.8
Fish		-	-	-	-	62	3.1
Unidentified		-	-	-	-	688	379.0
<i>Subtotal</i>		-	-	-	-	<i>1060</i>	<i>2145</i>
<b>Total</b>		<b>9</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>1164</b>	<b>3023.8</b>



Table 12. Number of vertebrate remains by phase from 41-49 Walmgate (Time Team site), York.

Taxon		Phase 1	Phase 3	Phase 4	Total no. frags
Vole/mouse	Microtine/Murine	1	-	-	1
Bank vole	<i>Clethrionomys glareolus</i> (Schreber)	-	-	1	1
Mouse	Murine sp.	-	1	-	1
Cat	<i>Felis f. domestic</i>	-	-	1	1
Pig	<i>Sus f. domestic</i>	1	1	14	16
Cattle	<i>Bos f. domestic</i>	5	-	19	24
Sheep/goat	Caprovid	1	-	11	12
Duck	<i>Anas sp.</i>	-	-	1	1
?Fowl	cf. <i>Gallus f. domestic</i>	-	-	1	1
Chicken	<i>Gallus f. domestic</i>	-	-	2	2
Herring	<i>Clupea harengus</i> L.	6	7	12	25
Eel	<i>Anguilla anguilla</i> (L.)	1	2	2	5
Cod family	Gadid	1	-	2	3
Cod	<i>Gadus morhua</i> L.	-	1	8	9
Haddock	<i>Melanogrammus aeglefinus</i> (L.)	-	-	2	2
<b>Subtotal</b>		<b>16</b>	<b>12</b>	<b>76</b>	<b>104</b>
Large mammal		22	6	98	126
Medium-sized mammal 1		18	19	127	164
Bird		6	2	12	20
Fish		4	19	39	62
Unidentified		132	105	451	688
<b>Subtotal</b>		<b>182</b>	<b>151</b>	<b>727</b>	<b>1060</b>
<b>Total</b>		<b>198</b>	<b>163</b>	<b>803</b>	<b>1164</b>

Table 13. Number of vertebrate remains by recovery method from 41-49 Walmgate (Time Team site), York.

Taxon		Hand-collected	SRS	BS	GBA	Total no. frags
Vole/mouse	Microtine/Murine	-	-	-	1	1
Bank vole	<i>Clethrionomys glareolus</i> (Schreber)	-	-	-	1	1
Mouse	Murine sp.	-	-	1	-	1
Cat	<i>Felis f. domestic</i>	1	-	-	-	1
Pig	<i>Sus f. domestic</i>	4	7	5	-	16
Cattle	<i>Bos f. domestic</i>	15	8	1	-	24
Sheep/goat	Caprovid	7	4	1	-	12
Duck	<i>Anas sp.</i>	1	-	-	-	1
?Fowl	cf. <i>Gallus f. domestic</i>	-	1	-	-	1
Chicken	<i>Gallus f. domestic</i>	1	1	-	-	2
Herring	<i>Clupea harengus L.</i>	-	-	19	6	25
Eel	<i>Anguilla anguilla (L.)</i>	-	-	3	2	5
Cod family	Gadid	-	1	1	1	3
Cod	<i>Gadus morhua L.</i>	-	8	1	-	9
Haddock	<i>Melanogrammus aeglefinus (L.)</i>	-	2	-	-	2
<b>Subtotal</b>		<b>29</b>	<b>32</b>	<b>32</b>	<b>11</b>	<b>104</b>
Large mammal		62	44	17	3	126
Medium-sized mammal 1		49	64	51	-	164
Bird		1	10	9	-	20
Fish		-	3	57	2	62
Unidentified		63	145	480	-	688
<b>Subtotal</b>		<b>175</b>	<b>266</b>	<b>614</b>	<b>5</b>	<b>1060</b>
<b>Total</b>		<b>204</b>	<b>298</b>	<b>646</b>	<b>16</b>	<b>1164</b>

Table 14. Proportions of meat-bearing and non meat-bearing parts of the skeleton, from 41-49 Walmgate (Time Team site), York

<b>Species</b>	<b>Meat-bearing</b>	<b>Non meat-bearing</b>	<b>Total</b>
Cattle	8	16	<b>24</b>
Caprovid	4	7	<b>11</b>
Pig	1	15	<b>16</b>
<b>Total</b>	<b>13</b>	<b>38</b>	<b>51</b>