

Table III.—Stature  $M_s = 67.2$  inches;  $Q_s = 1.75$  inches. Left Cubit  $M_c = 18.05$  inches;  $Q_c = 0.56$  inch.

No. of cases.	Stature.	Deviation from $M_s$ reckoned in		Mean of corresponding left cubits.	Deviation from $M_c$ reckoned in			Smoothed values multiplied by $Q_c$ .	Addition of $M_c$
		Inches.	Units of $Q_s$ .		Inches.	Units of $Q_c$			
						Observed.	Smoothed.		
30	70.0	+2.8	+1.69	18.8	+0.8	+1.42	+1.30	+0.73	18.05
50	69.0	+1.8	+1.03	18.3	+0.3	+0.53	+0.84	+0.47	18.05
38	68.0	+0.8	+0.46	18.2	+0.2	+0.36	+0.38	+0.21	18.05
61	67.0	-0.2	-0.11	18.1	+0.1	+0.18	-0.08	-0.04	18.05
48	66.0	-1.2	-0.69	17.8	-0.2	-0.36	-0.54	-0.30	17.25
36	65.0	-2.2	-1.25	17.7	-0.3	-0.53	-1.00	-0.56	17.25
21	64.0	-3.2	-1.83	17.2	-0.8	-1.46	-1.46	-0.80	17.25

  

No. of cases.	Left cubit.	Deviation from $M_c$ reckoned in		Mean of corresponding statures.	Deviation from $M_s$ reckoned in			Smoothed values multiplied by $Q_c$ .	Addition of $M_s$
		Inches.	Units of $Q_c$ .		Inches.	Units of $Q_s$			
						Observed.	Smoothed.		
38	19.25	+1.20	+2.14	70.3	+3.1	+1.8	+1.70	+3.0	70.3
55	18.75	+0.70	+1.25	68.7	+1.5	+0.9	+1.00	+1.8	69.25
102	18.25	+0.20	+0.36	67.4	+0.2	+0.1	+0.28	+0.5	67.4
61	17.75	-0.30	-0.53	66.3	-0.9	-0.5	-0.43	-0.8	66.3
49	17.25	-0.80	-1.42	65.0	-2.2	-1.3	-1.15	-2.0	65.0
25	16.75	-1.30	-2.31	63.7	-3.5	-2.0	-1.85	-3.2	64.0