

VALUES USED IN DERIVING DOUBLE-SAMPLING PLANS
WITH A SPECIFIED p_l AND p_2
($n_2 = 2n_1$, $\alpha = 0.05$, and $\beta = 0.10$)

Plan Number	R^*	Acceptance Numbers		Approximate Values of pn_l for			Approximate (ASN)/ n_l for 0.95 Point [†]
		c_1	c_2	$P = 0.95$	$P = 0.50$	$P = 0.10$	
1	14.33	0	1	0.16	0.84	2.32	1.299
2	7.80	0	2	0.31	1.07	2.43	1.535
3	6.52	1	3	0.60	1.80	3.90	1.243
4	5.42	0	3	0.49	1.35	2.64	1.770
5	5.11	1	4	0.77	1.97	3.93	1.360
6	4.30	0	4	0.68	1.64	2.93	1.987
7	4.18	1	5	0.96	2.19	4.01	1.498
8	4.11	2	6	1.30	2.89	5.34	1.285
9	3.67	0	5	0.89	1.95	3.26	2.177
10	3.61	2	7	1.49	3.06	5.37	1.377
11	3.58	1	6	1.16	2.44	4.17	1.648
12	3.27	0	6	1.11	2.26	3.62	2.340
13	3.26	3	9	2.06	3.97	6.70	1.307
14	3.22	2	8	1.69	3.28	5.45	1.482
15	2.98	3	10	2.26	4.14	6.74	1.385
16	2.92	2	9	1.91	3.52	5.58	1.598
17	2.81	4	12	2.85	5.03	8.02	1.321
18	2.75	3	11	2.47	4.35	6.81	1.473
19	2.70	2	10	2.14	3.78	5.76	1.720
20	2.63	4	13	3.06	5.21	8.06	1.390
21	2.57	3	12	2.70	4.58	6.93	1.570
22	2.53	2	11	2.37	4.06	6.00	1.845
23	2.48	4	14	3.28	5.41	8.13	1.467
24	2.42	3	13	2.93	4.84	7.09	1.673
25	2.35	4	15	3.51	5.64	8.23	1.552
26	2.30	3	14	3.17	5.11	7.30	1.780
27	2.24	4	16	3.74	5.89	8.39	1.642
28	2.21	3	15	3.41	5.39	7.55	1.889
29	2.15	4	17	3.98	6.16	8.58	1.736
30	2.14	3	16	3.66	5.69	7.83	1.997
31	2.08	4	18	4.23	6.44	8.81	1.834
32	2.02	4	19	4.49	6.72	9.06	1.931
33	1.97	4	20	4.75	7.02	9.35	2.028
34	1.93	4	21	5.01	7.33	9.65	2.122
35	1.89	4	22	5.27	7.64	9.97	2.213
36	1.82	5	25	6.12	8.66	11.15	2.146
37	1.72	6	30	7.52	10.30	12.96	2.249

* $R = p_2/p_1$.

[†] ASN is without curtailment on second sample.

VALUES USED IN DERIVING DOUBLE-SAMPLING PLANS
WITH A SPECIFIED p_l AND p_2
($n_2 = n_1$, $\alpha = 0.05$, and $\beta = 0.10$)

Plan Number	R^*	Acceptance Numbers		Approximate Values of pn_l for			Approximate (ASN)/ n_l for 0.95 Point [†]
		c_l	c_2	$P = 0.95$	$P = 0.50$	$P = 0.10$	
1	12.05	0	1	0.21	1.01	2.49	1.187
2	7.46	1	2	0.53	1.82	3.93	1.098
3	6.79	0	2	0.43	1.42	2.92	1.349
4	5.38	1	3	0.76	2.12	4.09	1.177
5	4.98	0	3	0.70	1.88	3.46	1.501
6	4.62	2	4	1.16	2.91	5.38	1.113
7	4.27	1	4	1.04	2.50	4.42	1.277
8	4.09	0	4	0.99	2.36	4.06	1.629
9	3.88	2	5	1.43	3.20	5.54	1.173
10	3.64	1	5	1.34	2.93	4.87	1.387
11	3.61	3	6	1.87	3.98	6.76	1.120
12	3.56	0	5	1.31	2.85	4.67	1.730
13	3.38	2	6	1.72	3.56	5.83	1.249
14	3.25	1	6	1.66	3.39	5.40	1.495
is	3.22	3	7	2.15	4.27	6.91	1.171
16	3.21	0	6	1.64	3.34	5.28	1.807
17	3.09	4	8	2.62	5.05	8.09	1.125
18	2.92	3	8	2.46	4.62	7.18	1.233
19	2.84	4	9	2.90	5.33	8.24	1.169
20	2.77	5	10	3.39	6.10	9.39	1.128
21	2.71	3	9	2.79	5.02	7.54	1.305
22	2.63	4	10	3.22	5.67	8.48	1.223
23	2.59	5	11	3.69	6.38	9.53	1.168
24	2.55	3	10	3.13	5.45	7.98	1.383
25	2.48	4	11	3.56	6.06	8.82	1.285
26	2.44	5	12	4.01	6.72	9.77	1.216
27	2.36	4	12	3.91	6.49	9.23	1.354
28	2.32	5	13	4.35	7.10	10.08	1.271
29	2.27	4	13	4.28	6.94	9.70	1.425
30	2.23	5	14	4.70	7.52	10.47	1.332
31	2.15	5	15	5.07	7.96	10.92	1.397
32	2.09	5	16	5.45	8.42	11.40	1.463

* $R = p_2/p_l$.

[†] ASN is without curtailment on second sample.