

# THE UNIVERSITY of York

# **Discussion Papers in Economics**

No. 2000/10

The Covariance Structure of Mixed ARMA Models

by

Menelaos Karanasos

Department of Economics and Related Studies University of York Heslington York, YO10 5DD

# THE COVARIANCE STRUCTURE OF MIXED ARMA MODELS

#### Menelaos Karanasos\*

University of York, Heslington, York, YO10 5DD, UK

#### Abstract

The purpose of this paper is to examine the covariance structure of mixed ARMA models, as discussed in Granger and Morris (1976). The method we use to obtain the autocovariances is based on the Wold representation of an ARMA model as it is given in Pandit (1973) or in Karanasos (2000). We give two examples to illustrate our general results: (i) two ARMA(2,2) processes with identical autoregressive polynomials and different moving average ones, and (ii) two ARMA(2,1) processes with different autoregressive and moving average polynomials.

Key Words: Autocovariance Structure, Mixed ARMA Models, Wold Representation. JEL Classification: C22

\* For correspondence: Email: mk16@york.ac.uk, Tel: 01904 433799, Fax: 01904 433759.

I would like to thank M. Karanassou and M. Dietrich for helpful comments and suggestions.

#### 1 Introduction

Ever since their introduction by Yule (1921, 1927) the autoregressive and moving average models have been greatly favoured by time series analysts. The purpose of this paper is to examine the covariance structure of mixed autoregressive and moving average (ARMA) time series models.

The theoretical autocovariance function (acf) of the ARMA model is an important tool in time series analysis. The calculation of the theoretical autocovariances is used (i) to estimate ARMA models with conventional exact maximum likelihood procedures (ii) to analyze the distribution of estimated ARMA parameters (Hannan, 1970) and (iii) to initialize simulations with ARMA models. The theoretical acf of the univariate ARMA(p,q) model has been already derived in the literature. The autocovariances are obtained in terms of the roots of the autoregressive (AR) polynomial and the parameters of the moving average (MA) one.

Pandit (1973, pp. 100, 141) and Pandit and Wu<sup>1</sup> (1983, pp. 105, 129-130), hereafter PW, derived the acf of the ARMA(p,q) model (when the roots of the AR polynomial are distinct) by using the infinite moving average (ima) or Wold representation<sup>2</sup> of the aforementioned model. Nerlove, Grether and Carvalho (1979, pp. 39, 78-85) used the canonical factorization (cf) of the autocovariance generating function (agf<sup>3</sup>) to derive a general expression for the acf of the ARMA(p,q) process (As an illustration they presented the acf of the MA(q), AR(p), and ARMA(1,1) models).

Zinde-Walsh (1988) obtained the acf of the ARMA(p,q) model by using the standard (see for example, Priestley, 1981<sup>4</sup>) spectral representation for the acf of a stationary ARMA process. She derived expressions for both cases of simple and multiple roots of the AR polynomial. Karanasos (1998), hereafter K, derived the acf of the ARMA(p,q) model (when the roots of the AR polynomial are distinct) by expressing the ARMA(p,q) model as an AR(1) process with an ARMA(p-1,q) error.

Algorithms for computing the theoretical autocovariances for univariate ARMA processes have been suggested in McLeod (1975, 1977) and Tunnicliffe Wilson (1979).

This paper contributes to the above literature by deriving the acf of the sum of ARMA processes when the roots of the AR polynomials are distinct. Granger and Morris (1976)

<sup>&</sup>lt;sup>1</sup>PW only examine the case where q < p.

<sup>&</sup>lt;sup>2</sup>They called the coefficient function in this expansion "Green's function" (see Miller (1968)). What PW have called Green's function is also referred to in the literature as weighting function (see Wiener (1949) and Pugachev (1957)) or as  $\psi$  weights (see Box and Jenkins (1970)).

For a proof of Wold's theorem see Wold (1938, pp. 75-89), Hannan (1970, pp.136-137), Anderson (1971, pp. 420-421), Sargent (1979, pp.257-262) or Brockwell and Davis (1987, pp. 180-182), hereafter BD; also see Hannan (1970, pp. 157-158) for the vector case. Wold's theorem has also been discussed by, for example, Nerlove, Grether and Carvalho (1979, pp. 30-36), hereafter NGC, PW (1983, pp. 87-89), and Reinsel (1993, p. 7), hereafter R.

The ima representation of the ARMA model has been discussed by, for example, O.D. Anderson (1976, p 44), PW (1983, p 108), Granger and Newbold (1986, pp. 25-26), hereafter GN, BD (1987, pp. 87-89), Wei (1989, p 56), Hamilton (1994, pp. 59-60), Gourieroux and Monfort (1997, pp 160-162); see also Mittnik (1987), R (1993, pp. 33-34), Lutkepohl (1993, pp. 220-221), hereafter L, or Gourieroux and Monfort (1997, p 252) for the vector case.

 $<sup>^{3}</sup>$ The cf of the agf has also been discussed by, for example, O.D. Anderson (1976, p 129), GN (1986, pp. 26-27), BD (1987, pp. 102-103), Wei (1989, pp. 242-243) or Hamilton (1994, pp. 61-63); see also BD (1987, p 410) or R(1993, pp. 33-34), for the vector case.

 $<sup>^{4}</sup>$ For the important subject of spectral analysis see also the excellent books by Jenkins and Watts (1968), and Hannan (1967).

consider such mixed ARMA models. As most economic series are both aggregates and are measured with error it follows that these mixed models are often found in practice. We examine three distinct cases, depending on whether the processes have identical or different autoregressive parts and uncorrelated or correlated errors. We obtain our results by using the ima representation of the ARMA(p,q) model, as it is given in Pandit (1973) or in K (2000)<sup>5</sup> of the ARMA(p,q) model. The autocovariances are expressed in terms of the roots of the AR polynomials and the parameters of the MA ones. In the case of distinct roots Zinde-Walsh (1988) or K(1998) results are special cases of our Theorem 1 (see Proposition 1a).

#### 2 MIXED ARMA MODELS

The following Proposition presents the autocovariances of an ARMA(p,q) model, when the roots of the AR polynomial are distinct.

Let  $y_t$  be an ARMA(p,q) process given by:

$$\Phi(L)y_t = \mu + \Theta(L)\epsilon_t, \ \epsilon_t \sim IID(0, \sigma_\epsilon^2), \tag{2.1}$$

$$\Theta(L) = \sum_{k=0}^{q} \theta_k L^k, \ \Phi(L) = -\sum_{k=0}^{p} \phi'_k L^k = \prod_{k=1}^{p} (1 - \phi_k L), \ \phi'_0 = 1, \ \phi_k \neq \phi_l$$
(2.1*a*)

Assumption 1. All the roots of the autoregressive polynomial  $[\Phi(L)]$  lie outside the unit circle (Stationarity condition).

Assumption 2. The polynomials  $\Phi(L)$  and  $\Theta(L)$  are left coprime. In other words the representation  $\frac{\Phi(L)}{\Theta(L)}$  is irreducible.

Proposition 1a. Under assumptions 1-2 the autocovariances of  $y_t$  ( $\gamma_m$ ) are given by<sup>6</sup>

$$\gamma_m = \sum_{l=1}^p \zeta_{lm} \lambda_{l,min(m,q)} \sigma_{\epsilon}^2, \ \zeta_{lm} = \frac{\phi_l^{p-1+m}}{\prod_{k=1}^p (1-\phi_l \phi_k) \prod_{\substack{k=1\\k \neq l}}^p (\phi_l - \phi_k)},$$
(2.2)

$$\lambda_{lm} = \sum_{k=0}^{q} \theta_k^2 + \sum_{d=1}^{m} \sum_{k=0}^{q-d} \theta_k \theta_{k+d} (\phi_l^d + \phi_l^{-d}) + \sum_{d=m+1}^{q} \sum_{k=0}^{q-d} \theta_k \theta_{k+d} (\phi_l^d + \phi_l^{d-2m})$$
(2.2a)

The proof of Proposition 1a is given in Appendix A.

By considering the model generating the sum of two or more series, Granger and Morris (1976) have shown that the mixed ARMA model is the one most likely to occur.

 $<sup>{}^{5}</sup>$ K has also used the ima representation in the context of the GARCH and GARCH in mean models.

<sup>&</sup>lt;sup>6</sup>Zinde Victoria Walsh (1988) derived the above result (eqs 2a and 2b) by using the standard (see for example, Priestley (1981)) spectral representation for the acf of a stationary ARMA process whereas K(1998) derived the same result by expressing the ARMA(p,q) model as an AR(1) process with an ARMA(p-1,q) error. Zinde-Walsh's formula is general as it is not restricted to the case of simple roots.

They mentioned that "two situations where series are added together are of particular interpretational importance. The first is where series are aggregated to form some total, and the second one is where the observed series is the sum of the true process plus an observational error. Most of the macroeconomic series, such as GNP, unemployment or exports are aggregates. Virtually any macroeconomic series, other than certain prices or interest rates, contain important observation errors." So speaking the following analysis of the autocovariance function of the sum of ARMA processes can be a useful tool for the applied economist.

Let  $y_t$  be equal to the sum of n uncorrelated ARMA $(p, q_i)$  processes that have identical autoregressive parts:

$$y_{t} = \sum_{i=1}^{n} y_{it}, \ \Phi(L)y_{it} = \Theta_{i}(L)\epsilon_{it}, \ \Theta_{i}(L) = \sum_{k=0}^{q_{i}} \theta_{ik}L^{k},$$
(2.3)

$$\bar{\epsilon}_t = \begin{bmatrix} \epsilon_{1t} \\ \cdot \\ \cdot \\ \cdot \\ \epsilon_{nt} \end{bmatrix}, \ \bar{\epsilon}_t \sim IID(0, \bar{\sigma}_\epsilon^2), \ \bar{\sigma}_\epsilon^2 = \begin{bmatrix} \sigma_{11,\epsilon} & \dots & 0 \\ \cdots & \cdots & \cdots \\ 0 & \dots & \sigma_{nn,\epsilon} \end{bmatrix}$$
(2.3a)

where  $\Phi(L)$  is defined by (2.1*a*).

Assumption 3. The polynomials  $\Phi(L)$  and  $\Theta_i(L)$  are left coprime. In other words the representation  $\frac{\Phi(L)}{\Theta_i(L)}$  is irreducible.

Proposition  $\hat{1b}$ . Under assumptions 1 and 3 the autocovariances of the preceding ARMA process are given by

$$\gamma_m = \sum_{i=1}^n \sum_{l=1}^p \zeta_{lm} \lambda^i_{l,min(m,q_i)} \sigma_{ii,\epsilon},$$
(2.4)

$$\lambda_{lm}^{i} = \sum_{d=0}^{q_{i}} \theta_{id}^{2} + \sum_{d=1}^{m} \sum_{r=0}^{q_{i}-d} \theta_{ir} \theta_{ir+d} (\phi_{l}^{d} + \phi_{l}^{-d}) + \sum_{d=m+1}^{q_{i}} \sum_{r=0}^{q_{i}-d} \theta_{ir} \theta_{ir+d} (\phi_{l}^{d} + \phi_{l}^{d-2m})$$
(2.4*a*)

where the  $\zeta_{lm}$  is defined in (2.2).

The Proof of Proposition 1b is given in Appendix A.

In the Proposition that follows the assumption of independence is weakened to allow for contemporaneous correlation between the noise series.

Let  $y_t$  be equal to the sum of n correlated ARMA $(p, q_i)$  processes as defined by (2.3), where  $\bar{\sigma}_{\epsilon}^2$  is given by

$$\bar{\sigma}_{\epsilon}^{2} = \begin{bmatrix} \sigma_{11,\epsilon} & \dots & \sigma_{1n,\epsilon} \\ \dots & \dots \\ \vdots \\ \sigma_{n1,\epsilon} & \dots & \sigma_{nn,\epsilon} \end{bmatrix}$$
(2.5)

Proposition 1c. Under assumptions 1 and 3 the autocovariances of  $y_t$  are given by

$$\gamma_m = \sum_{j=1}^n \sum_{i=1}^n \sum_{l=1}^p \zeta_{lm} \lambda_{l,m}^{ij} \sigma_{ij,\epsilon}, \qquad (2.6)$$

$$\lambda_{l,m}^{ij} = \sum_{c=0}^{q_j} \sum_{d=0}^{q'_i} \theta_{id} \theta_{j,d+c} \phi_l^c + \sum_{c=1}^{m\star} \sum_{d=0}^{q'_j} \theta_{jd} \theta_{id+c} \phi_l^{-c} + \sum_{c=m^\star+1}^{q_i} \sum_{d=0}^{q'_j} \theta_{jd} \theta_{id+c} \phi_l^{c-2m}$$
(2.6a)

where

$$q'_{i} = min(q_{i}, q_{j} - c) \qquad q'_{j} = min(q_{j}, q_{i} - c)$$
  
$$m^{\star} = min(m, q_{i}) \qquad (2.6b)$$

and  $\zeta_{lm}$  is defined in (2.2). The proof of Proposition 1c is given in Appendix A.

*Example 1.* Consider two ARMA(2,2) processes  $(y_{1t}, y_{2t})$  with identical autoregressive polynomials and different moving average ones

$$(1 - \phi_1 L)(1 - \phi_2 L)y_{1t} = (1 + \theta_{11}L + \theta_{12}L^2)\epsilon_{1t}, \quad \phi_1 \neq \phi_2, \quad |\phi_1|, |\phi_2| < 1$$
$$(1 - \phi_1 L)(1 - \phi_2 L)y_{2t} = (1 + \theta_{21}L + \theta_{22}L^2)\epsilon_{2t},$$
$$\begin{bmatrix} \epsilon_{1t} \\ \epsilon_{2t} \end{bmatrix} \sim IID(0, \bar{\sigma}_{\epsilon}^2), \quad \bar{\sigma}_{\epsilon}^2 = \begin{bmatrix} var(\epsilon_{1t}) & cov(\epsilon_{1t}, \epsilon_{2t}) \\ cov(\epsilon_{1t}, \epsilon_{2t}) & var(\epsilon_{2t}) \end{bmatrix} = \begin{bmatrix} \sigma_{11,\epsilon} & \sigma_{12,\epsilon} \\ \sigma_{12,\epsilon} & \sigma_{22,\epsilon} \end{bmatrix}$$

The cross covariances between the two processes are

$$\begin{aligned} \cos(y_{1t}, y_{2t}) &= \sigma_{12,\epsilon} \times \\ &\{ \frac{\phi_1 [1 + \theta_{11} \theta_{21} + \theta_{12} \theta_{22} + (\theta_{21} + \theta_{11} \theta_{22} + \theta_{11} + \theta_{21} \theta_{12}) \phi_1 + (\theta_{12} + \theta_{22}) \phi_1^2]}{(1 - \phi_1 \phi_2) (1 - \phi_1^2) (\phi_1 - \phi_2)} &+ \\ &\frac{\phi_2 [1 + \theta_{11} \theta_{21} + \theta_{12} \theta_{22} + (\theta_{21} + \theta_{11} \theta_{22} + \theta_{11} + \theta_{21} \theta_{12}) \phi_2 + (\theta_{12} + \theta_{22}) \phi_2^2]}{(1 - \phi_1 \phi_2) (1 - \phi_2^2) (\phi_2 - \phi_1)} \end{aligned}$$

 $cov(y_{1t}, y_{2,t-1}) = \sigma_{12,\epsilon} \times$ 

$$\{ \frac{\phi_1^2 [1 + \theta_{11}\theta_{21} + \theta_{12}\theta_{22} + (\theta_{21} + \theta_{11}\theta_{22})\phi_1 + (\theta_{11} + \theta_{21}\theta_{12})\phi_1^{-1} + \theta_{22}\phi_1^2 + \theta_{12}\phi_1^0]}{(1 - \phi_1\phi_2)(1 - \phi_1^2)(\phi_1 - \phi_2)} \\ + \frac{\phi_2^2 [1 + \theta_{11}\theta_{21} + \theta_{12}\theta_{22} + (\theta_{21} + \theta_{11}\theta_{22})\phi_2 + (\theta_{11} + \theta_{21}\theta_{12})\phi_2^{-1} + \theta_{22}\phi_2^2 + \theta_{12}\phi_2^0]}{(1 - \phi_1\phi_2)(1 - \phi_2^2)(\phi_2 - \phi_1)} \}$$

 $cov(y_{2t}, y_{1,t-1}) = \sigma_{12,\epsilon} \times$ 

$$\{ \frac{\phi_1^2 [1 + \theta_{11}\theta_{21} + \theta_{12}\theta_{22} + (\theta_{11} + \theta_{21}\theta_{12})\phi_1 + (\theta_{21} + \theta_{11}\theta_{22})\phi_1^{-1} + \theta_{12}\phi_1^2 + \theta_{22}\phi_1^0]}{(1 - \phi_1\phi_2)(1 - \phi_1^2)(\phi_1 - \phi_2)} \\ + \frac{\phi_2^2 [1 + \theta_{11}\theta_{21} + \theta_{12}\theta_{22} + (\theta_{11} + \theta_{21}\theta_{12})\phi_2 + (\theta_{21} + \theta_{11}\theta_{22})\phi_2^{-1} + \theta_{12}\phi_2^2 + \theta_{22}\phi_2^0]}{(1 - \phi_1\phi_2)(1 - \phi_2^2)(\phi_2 - \phi_1)} \}$$

 $cov(y_{1t}, y_{2,t-\underset{m\geq 2}{m}}) = \sigma_{12,\epsilon} \times$ 

$$\left\{\frac{\phi_1^{1+m}[1+\theta_{11}\theta_{21}+\theta_{12}\theta_{22}+(\theta_{21}+\theta_{11}\theta_{22})\phi_1+(\theta_{11}+\theta_{21}\theta_{12})\phi_1^{-1}+\theta_{22}\phi_1^2+\theta_{12}\phi_1^{-2}]}{(1-\phi_1\phi_2)(1-\phi_1^2)(\phi_1-\phi_2)}\right\}$$

$$+\frac{\phi_{2}^{1+m}[1+\theta_{11}\theta_{21}+\theta_{12}\theta_{22}+(\theta_{21}+\theta_{11}\theta_{22})\phi_{2}+(\theta_{11}+\theta_{21}\theta_{12})\phi_{2}^{-1}+\theta_{22}\phi_{2}^{2}+\theta_{12}\phi_{2}^{-2}]}{(1-\phi_{1}\phi_{2})(1-\phi_{2}^{2})(\phi_{2}-\phi_{1})}\}$$

$$cov(y_{2t}, y_{1,t-\underset{m\geq 2}{m\geq 2}}) = \sigma_{12,\epsilon} \times$$

$$\{\frac{\phi_1^{1+m}[1+\theta_{11}\theta_{21}+\theta_{12}\theta_{22}+(\theta_{11}+\theta_{21}\theta_{12})\phi_1+(\theta_{21}+\theta_{11}\theta_{22})\phi_1^{-1}+\theta_{12}\phi_1^2+\theta_{22}\phi_1^{-2}]}{(1-\phi_1\phi_2)(1-\phi_1^2)(\phi_1-\phi_2)}$$

$$+\frac{\phi_{2}^{1+m}[1+\theta_{11}\theta_{21}+\theta_{12}\theta_{22}+(\theta_{11}+\theta_{21}\theta_{12})\phi_{2}+(\theta_{21}+\theta_{11}\theta_{22})\phi_{2}^{-1}+\theta_{12}\phi_{2}^{2}+\theta_{22}\phi_{2}^{-2}]}{(1-\phi_{1}\phi_{2})(1-\phi_{2}^{2})(\phi_{2}-\phi_{1})}\}$$

Note, that when the two moving average polynomials and the two error terms are identical  $(y_{1t} = y_{2t} = y_t)$  the above expressions give the autocovariances of the  $y_t$  process:  $cov(y_{1t}, y_{2,t-m}) = cov(y_{2t}, y_{1,t-m}) = cov_m(y_t).$ 

In the following Theorem the assumption of identical autoregressive parts is relaxed.

Let  $y_t$  be equal to the sum of n correlated ARMA $(p_i, q_i)$  processes, where the roots of the AR polynomials are distinct:

$$y_t = \sum_{i=1}^n y_{it}, \ \Phi_i(L)y_{it} = \Theta_i(L)\epsilon_{it}, \ \Phi_i(L) = -\sum_{k=0}^{p_i} \phi'_{ik}L^k = \prod_{k=1}^{p_i} (1 - \phi_{ik}L), \ \phi_{ik} \neq \phi_{il}$$
(2.7)

where  $\Theta_i(L)$ ,  $\bar{\epsilon}_t$ , are defined by (2.3) and  $\bar{\sigma}_{\epsilon}^2$  is defined by (2.5).

Assumption 4. All the roots of the autoregressive polynomials  $[\Phi_i(L)]$  lie outside the unit circle (Stationarity conditions).

Assumption 5. The polynomials  $\Phi_i(L)$  and  $\Theta_i(L)$  are left coprime. In other words the representation  $\frac{\Phi_i(L)}{\Theta_i(L)}$  is irreducible.

Theorem 1. Under assumptions 4 and 5 the autocovariances of the preceding process are given by

$$\gamma_m = \sum_{j=1}^n \sum_{i=1}^n \left( \sum_{l=1}^{p_i} \zeta_{il,j}^m \lambda_{l,m}^{ij} + \sum_{k=1}^{p_j} \zeta_{jk,i}^m \lambda_{ij}^{k,m} \right) \sigma_{ij,\epsilon},$$
(2.8)

$$\zeta_{il,j}^{m} = \frac{\zeta_{il}^{m}}{\prod_{k=1}^{p_{j}} (1 - \phi_{il}\phi_{jk})}, \quad \zeta_{il}^{m} = \frac{\phi_{il}^{p_{i}-1+m}}{\prod_{\substack{k=1\\k \neq l}}^{p_{i}} (\phi_{il} - \phi_{ik})}, \quad (2.8a)$$

$$\lambda_{l,m}^{ij} = \sum_{c=0}^{q_j} \sum_{d=0}^{q'_i} \theta_{id} \theta_{j,d+c} \phi_{il}^c + \sum_{c=1}^{m^*} \sum_{d=0}^{q'_j} \theta_{jd} \theta_{id+c} \phi_{il}^{-c}, \quad \lambda_{ij}^{k,m} = \sum_{c=m^*+1}^{q_i} \sum_{d=0}^{q'_j} \theta_{jd} \theta_{id+c} \phi_{jk}^{c-2m}$$
(2.8b)

The proof of Theorem 1 is given in Appendix B.

*Example 2.* Consider two ARMA(2,1) processes  $(y_{1t}, y_{2t})$  with different autoregressive and moving average polynomials

$$(1 - \phi_{11}L)(1 - \phi_{12}L)y_{1t} = (1 + \theta_{11}L)\epsilon_{1t}, \quad \phi_{11} \neq \phi_{12}, \quad |\phi_{11}|, |\phi_{12}| < 1$$
$$(1 - \phi_{21}L)(1 - \phi_{22}L)y_{2t} = (1 + \theta_{21}L)\epsilon_{2t}, \quad \phi_{21} \neq \phi_{22}, \quad |\phi_{21}|, |\phi_{22}| < 1$$
$$\begin{bmatrix} \epsilon_{1t} \\ \epsilon_{2t} \end{bmatrix} \sim IID(0, \bar{\sigma}_{\epsilon}^{2}), \quad \bar{\sigma}_{\epsilon}^{2} = \begin{bmatrix} var(\epsilon_{1t}) & cov(\epsilon_{1t}, \epsilon_{2t}) \\ cov(\epsilon_{1t}, \epsilon_{2t}) & var(\epsilon_{2t}) \end{bmatrix} = \begin{bmatrix} \sigma_{11,\epsilon} & \sigma_{12,\epsilon} \\ \sigma_{12,\epsilon} & \sigma_{22,\epsilon} \end{bmatrix}$$

The cross covariances between the two processes are

 $cov(y_{1t}, y_{2t}) = \sigma_{12,\epsilon} \times$ 

$$\left\{\frac{\phi_{11}[1+\theta_{11}\theta_{21}+\theta_{21}\phi_{11}]}{(\phi_{11}-\phi_{12})(1-\phi_{11}\phi_{21})(1-\phi_{11}\phi_{22})}+\frac{\phi_{12}[1+\theta_{11}\theta_{21}+\theta_{21}\phi_{12}]}{(\phi_{12}-\phi_{11})(1-\phi_{12}\phi_{21})(1-\phi_{12}\phi_{22})}+\right.$$

$$+ \frac{\phi_{21}^2 \theta_{11}}{(\phi_{21} - \phi_{22})(1 - \phi_{21}\phi_{11})(1 - \phi_{21}\phi_{12})} + \frac{\phi_{22}^2 \theta_{11}}{(\phi_{22} - \phi_{21})(1 - \phi_{22}\phi_{11})(1 - \phi_{22}\phi_{12})} \}$$

$$cov(y_{1t}, y_{2,t-m}) = \{ \frac{\phi_{11}^{1+m}[1 + \theta_{11}\theta_{21} + \theta_{21}\phi_{11} + \theta_{11}\phi_{11}^{-1}]}{(\phi_{11} - \phi_{12})(1 - \phi_{11}\phi_{21})(1 - \phi_{11}\phi_{22})} + \frac{\phi_{12}^{1+m}[1 + \theta_{11}\theta_{21} + \theta_{21}\phi_{12} + \theta_{11}\phi_{12}^{-1}]}{(\phi_{12} - \phi_{11})(1 - \phi_{12}\phi_{21})(1 - \phi_{12}\phi_{22})} \} \sigma_{12,\epsilon}, \ m \ge 1$$

$$cov(y_{2t}, y_{1,t-m}) = \left\{ \frac{\phi_{21}^{1+m} [1 + \theta_{11}\theta_{21} + \theta_{11}\phi_{21} + \theta_{21}\phi_{21}^{-1}]}{(\phi_{21} - \phi_{22})(1 - \phi_{21}\phi_{11})(1 - \phi_{21}\phi_{12})} + \right\}$$

$$+\frac{\phi_{22}^{1+m}[1+\theta_{11}\theta_{21}+\theta_{11}\phi_{22}+\theta_{21}\phi_{22}^{-1}]}{(\phi_{22}-\phi_{21})(1-\phi_{22}\phi_{11})(1-\phi_{22}\phi_{12})}\}\sigma_{12,\epsilon}, \ m \ge 1$$

Note that when the autoregressive polynomials of the two processes are identical ( $\phi_{21} = \phi_{11}, \phi_{22} = \phi_{12}$ ) the above expressions reduce to those given in Example 1 with  $\theta_{12} = \theta_{22} = 0$ .

#### 3 Concluding Remarks

Since the publication of the book by Box and Jenkins, almost 30 years ago, the interest by time series analysts and econometricians in more complicated even if more parsimonious linear time series models has greatly increased. The purpose of this paper was to examine the covariance structure of mixed ARMA models. In Section 2 we presented the autocovariance function of the sum of n ARMA $(p_i, q_i)$  processes using the infinite moving average representation of an ARMA model. The autocovariances were expressed in terms of the roots of the AR polynomials and the parameters of the MA polynomials. In this paper we examined only the case of distinct roots of the autoregressive polynomials-the case of equal roots is left for future research. The ima representation technique can also be applied to non linear multivariate time series models (e.g. GARCH or Markov Switching) to derive their covariance structure<sup>7</sup>.

#### REFERENCES

Anderson, O. D. (1976) Time series analysis and forecasting: the Box-Jenkins approach, London: Butterworths.

Anderson, T. W. (1971) The statistical analysis of time series, New York: Wiley

Box, G.E.P., & G.M. Jenkins (1970) *Time series analysis forecasting and control*, Rev. Ed. 1976, San Francisco: Holden Day.

Brockwell, P. J., & R. A. Davis (1987) *Time series: theory and methods*, New York: Springer-Verlag.

Gourieroux, C. & A. Monfort (1997) *Time Series and Dynamic Models*, Cambridge: Cambridge University Press.

Granger, C.W.J., & M.J. Morris (1976) Time series modelling and interpretation, Journal of Royal Statistical Society 139, 246-257.

Granger, C.W.J., & P. Newbold (1986) Forecasting Economic Time Series, 2nd ed. London: Academic Press.

Hamilton, J.D. (1994) *Time Series Analysis*, Princeton: Princeton University Press. Hannan, E. J. (1967) *Time series analysis*, London: Science Paperbacks.

 $<sup>^{7}</sup>$ K (1999) has applied these techniques in order to obtain the covariance structure of various univariate and multivariate GARCH models.

Hannan, E.J. (1970) Multiple Time Series, New York: Wiley.

Jenkins, G. M., & D. G. Watts, (1968), *Spectral analysis and its applications*, San Francisco: Holden-Day.

Karanasos, M. (1998) A New method for obtaining the autocovariance of an ARMA model: an exact form solution, *Econometric Theory*, 14, 622-640.

Karanasos, M. (1999) The covariance structure of the S-GARCH and M-GARCH models, Unpublished mimeo, University of York.

Karanasos, M. (2000) Prediction in ARMA models with GARCH in mean effects. forthcoming in Journal of Time Series Analysis.

Lutkepohl, H. (1993) Introduction to multiple time series analysis, Berlin: Springer-Verlag.

Mc Leod, I. (1975) Derivation of the theoretical autocovariance function of autoregressivemoving average time series, *Applied Statistics* 24, 255-256, correction, 26 (1977), 194.

Miller, K. S. (1968) *Linear Difference Equations*, New York: W. A. Benjamin.

Mittnik, S. (1987) Non-recursive methods for computing the coefficients of the autoregressive and the moving-average representation of mixed ARMA processes, *Economic Letters*, 23, 279-284.

Nerlove, M., Grether, D. M., & J. L. Carvalho (1979) Analysis of economic time series: a synthesis, New York: Academic Press.

Pandit, S. M. (1973) Data dependent systems: modeling analysis and optimal control via time series, Ph.D. Thesis, University of Wiskonsin-Madison.

Pandit, S. M., & S. M. Wu (1983) *Time series and system analysis with applications*, New York: Wiley.

Priestley, M. B. (1981) Spectral analysis and time series, London: Academic Press.

Pugachev, V. S. (1957) Theory of random functions and its applications to automatic control problems, (Russian) English Trans. of 3rd Ed. by O. M. Blum, New York: Pargamon Press, 1965.

Reinsel, G. C. (1993) *Elements of multivariate time series analysis*, New York: Springer-Verlag.

Sargent, T. J. (1979) *Macroeconomic Theory*, New York: Academic Press, (2nd Ed. 1987).

Tunnicliffe Wilson, G. (1979) Some efficient computational procedures for high order ARMA models, *Journal of Statistical Computation and Simulation*, 8, 301-309.

Wei, W. S. (1989) *Time series analysis: univariate and multivariate methods*, New York: Addison Wesley.

Wiener, N. (1949) Extrapolation, interpolation and smoothing of stationary time series, New York: Wiley.

Wold, H. O. (1938) A study in the analysis of stationary time series, Stockholm: Almquist and Wicksell, (2nd Ed. 1954).

Yule, G.U. (1921) On the time correlation problem, with especial reference to the variate-difference correlation method, *Journal of Royal Statistical Society*, Series A, 84, 497-526.

Yule, G.U. (1927) On a method of investigating periodicities in disturbed series, with special reference to Wolfer's sunspot numbers, *Philosophical Transactions*, A226, p. 267.

Zinde-Walsh, V. (1988) Some exact formulae for autoregressive moving average processes, *Econometric Theory*, 4, 384-402.

# Appendix

## A PROOF OF PROPOSITION 1a

The impulse response function (IRF) of  $y_t$  is given by<sup>8</sup>

$$y_{t} = \sum_{r=0}^{\infty} e^{r} \epsilon_{t-r}, \text{ where } e^{r} = \sum_{l=1}^{p} \sum_{k=0}^{\min(r,q)} \zeta_{l}^{r-k} \theta_{k}, \quad \zeta_{l}^{m} = \frac{\phi_{l}^{p-1+m}}{\prod\limits_{\substack{k=1\\k \neq l}}^{p} (\phi_{l} - \phi_{k})}$$
(A.1)

From the preceding equation we have that

$$\gamma_m = cov_m(y_t) = \sum_{r=0}^{\infty} e^r e^{r+m} \sigma_{\epsilon}^2$$
(A.1*a*)

After some algebra we get

$$\gamma_m = \sum_{l=1}^p \zeta_l^m s_l^0 \lambda_{lm} \sigma_\epsilon^2, \text{ where } s_l^0 = \sum_{k=1}^p \frac{\zeta_k^0}{(1 - \phi_l \phi_k)}$$
(A.1b)

Subsequently, using

$$s_l^0 = \frac{1}{\prod_{k=1}^p (1 - \phi_l \phi_k)}$$
(A.1c)

we get (2.2)-(2.2a).

## PROOF OF PROPOSITION 1b

From (2.3) and using Proposition 1a we get

$$\gamma_m = \sum_{i=1}^n cov_m(y_{it}) = \sum_{i=1}^n \sum_{l=1}^p \zeta_{lm} \lambda_{l,min(m,q_i)}^i \sigma_{ii,\epsilon}$$
(A.2)

 $<sup>^{8}</sup>$ It is given in Pandit (1973) or K (2000).

## PROOF OF PROPOSITION 1c

The IRF of  $y_{it}$  is given by

$$y_{it} = \sum_{r=0}^{\infty} e_i^r \epsilon_{it-r}, \text{ where } e_i^r = \sum_{l=1}^p \sum_{k=0}^{\min(r,q_i)} \zeta_l^{r-k} \theta_{ik}$$
 (A.3)

From the preceding equation we get

$$cov(y_{it}, y_{jt-m}) = \sum_{r=0}^{\infty} e_j^r e_i^{r+m} \sigma_{ij,\epsilon}$$
(A.3*a*)

After some algebra we have that

$$cov(y_{it}, y_{jt-m}) = \sum_{l=1}^{p} \zeta_m^l s_l^0 \lambda_{l,m}^{ij} \sigma_{ij,\epsilon} = \sum_{l=1}^{p} \zeta_{lm} \lambda_{l,m}^{ij} \sigma_{ij,\epsilon}$$
(A.3b)

Using the preceding equation, together with equations (2.3) and (2.5) we get

$$\gamma_m = \sum_{j=1}^n \sum_{i=1}^n cov(y_{it}, y_{jt-m}) = \sum_{j=1}^n \sum_{i=1}^n \sum_{l=1}^p \zeta_{lm} \zeta_{l,m}^{ij} \sigma_{ij,\epsilon}$$
(A.3c)

## B PROOF OF THEOREM 1

The IRF of  $y_{it}$  is given by

$$y_{it} = \sum_{r=0}^{\infty} e_i^r \epsilon_{it-r}, \text{ where } e_i^r = \sum_{l=1}^{p_i} \sum_{k=0}^{\min(r,q_i)} \zeta_{il}^{r-k} \theta_{ik}, \quad \zeta_{il}^m = \frac{\phi_{il}^{p_i-1+m}}{\prod\limits_{\substack{k=1\\k \neq l}}^{p_i} (\phi_{il} - \phi_{ik})}$$
(B.1)

From the preceding equation we have

$$cov(y_{it}, y_{j,t-m}) = \sum_{r=0}^{\infty} e_j^r e_i^{r+m} \sigma_{ij,\epsilon}$$
(B.1a)

After some algebra we get

$$cov(y_{it}, y_{j,t-m}) = \left(\sum_{l=1}^{p_i} \zeta_{il}^m s_{j,il}^0 \lambda_{l,m}^{ij} + \sum_{k=1}^{p_j} \zeta_{jk}^m s_{i,jk}^0 \lambda_{ij}^{km}\right) \sigma_{ij,\epsilon}, \text{ where } s_{j,il}^0 = \sum_{k=1}^{p_j} \zeta_{jk}^{0,il},$$
(B.1b)

$$\zeta_{jk}^{0,il} = \frac{\zeta_{jk}^{0}}{(1 - \phi_{il}\phi_{jk})}, \quad \lambda_{l,m}^{ij} = \sum_{c=0}^{q_j} \sum_{d=0}^{q'_i} \theta_{id}\theta_{j,d+c}\phi_{il}^c + \sum_{c=1}^{m^*} \sum_{d=0}^{q'_j} \theta_{jd}\theta_{i,d+c}\phi_{il}^{-c} \tag{B.1c}$$

$$\lambda_{ij}^{km} = \sum_{c=m+1}^{q_i} \sum_{d=0}^{q'_j} \theta_{jd} \theta_{id+c} \phi_{jk}^{c-2m}, \tag{B.1d}$$

and  $q'_i = min(q_i, q_j - c), q'_j = min(q_j, q_i - c), m^* = min(m, q_i).$ In the preceding equation, we use

$$s_{j,il}^{0} = \frac{1}{\prod_{k=1}^{p_{j}} (1 - \phi_{il}\phi_{jk})}$$
(B.1e)

to get

$$cov(y_{it}, y_{j,t-m}) = \left(\sum_{l=1}^{p_i} \zeta_{il,j}^m \lambda_{l,m}^{ij} + \sum_{k=1}^{p_j} \zeta_{jk,i}^m \lambda_{ij}^{km}\right) \sigma_{ij,\epsilon}, \text{ where } \zeta_{il,j}^m = \frac{\zeta_{il}^m}{\prod_{k=1}^{p_j} (1 - \phi_{il}\phi_{jk})}$$
(B.1*f*)

Thus, we have

$$y_{t} = \sum_{i=1}^{n} y_{it} \Rightarrow cov_{m}(y_{t}) = \sum_{j=1}^{n} \sum_{i=1}^{n} cov(y_{it}, y_{j,t-m})$$
$$= \sum_{j=1}^{n} \sum_{i=1}^{n} (\sum_{l=1}^{p_{i}} \zeta_{il,j}^{m} \lambda_{l,m}^{ij} + \sum_{k=1}^{p_{j}} \zeta_{jk,i}^{m} \lambda_{ij}^{km}) \sigma_{ij,\epsilon}$$
(B.1g)