

HEDG

HEALTH, ECONOMETRICS AND DATA GROUP

THE UNIVERSITY *of York*

WP 15/05

Vertical Transmission of Overweight: Evidence from English Adoptees

Joan Costa-Font, Mireia Jofre-Bonet & Julian Le Grand

May 2015

<http://www.york.ac.uk/economics/postgrad/herc/hedg/wps/>

Vertical Transmission of Overweight: Evidence from English Adoptees

Joan Costa-Font^a, Mireia Jofre-Bonet^b, and Julian Le Grand^a

^a London School of Economics and Political Science (LSE), London, UK

^b Department of Economics, City University London, UK, and LSE Health, LSE.

Corresponding author: Mireia Jofre-Bonet, Department of Economics City University London, EC1V 0HB London, UK. Email: mireia.jofre-bonet@city.ac.uk . Other authors' emails: j.costa-font@lse.ac.uk; j.legrand@lse.ac.uk.

Acknowledgements: We would like to thank Richard Layard, Alice Mesnard, Alistair McGuire, Nick Powdthavee, Jan De Neuve, Nele Warrier, Berkay Ozcan, Mauro Laudicella, Marcos Vera-Hernandez, Brendan Walsh, Victoria Serra-Sastre, Patricia Cubi-Molla, and other participants in the LHEG at Kings College and the CEP Wellbeing Seminar Series at the LSE.

Abstract

We examine the vertical transmission of overweight drawing upon a sample of English children, both adopted and non-adopted, and their families. Our results suggest strong evidence of an intergenerational association of overweight among adoptees, indicating transmission through cultural factors. We find that, when both adoptive parents are overweight, the likelihood of an adopted child being overweight is between 10% and 20% higher than when they are not. We also find that the cultural transmission of overweight is not aggravated by having a full-time working mother, so do not confirm the existence of a female labour market participation penalty on child overweight among adoptees. Overall, our findings, despite subject to data limitations, are robust to a battery of robustness checks, specification and sample selection corrections.

Keywords: vertical transmission, cultural transmission, overweight, children, natural parents, Body Mass Index, sample selection.

JEL: I18, D13, Z1.

1. Introduction

Overweight and obesity, as a form of extreme overweight in children is of growing concern.

Evidence from the Health Survey for England suggests that the prevalence of overweight among 2-10 (11-15) year-olds averaged over the three years 2010 to 2012 was as high as 26% (35%), and obesity 13% (9%).¹ Nor is the situation any better in other parts of the United Kingdom (UK).² Even more concerning, estimates from the International Association for the Study of Obesity (IASO, 2011) indicate that the rates of overweight (including obese) children aged 5-17 years in the UK are among the highest in Europe and have experienced an increasing trend in the last decade, with a corresponding associated rising burden of morbidity (Berenson et al, 1993).

The mechanisms contributing to what might fairly be described as a *childhood overweight epidemic* are contentious, as are the appropriate policy interventions. A major problem for policy intervention is the identification of the relative importance of hereditary factors and environmental ones.

Childhood obesity is found to be partly heritable in studies of identical twins, but the estimates vary from 37 to 90% (Llewellyn 2003). Although we do not attempt to provide a comprehensive review of the growing literature on transmission of obesity, recent estimates using adoptees vary from 20 to 60% (Elks et al, 2012). In contrast, overweight in children seems to be significantly more influenced by the specific individual cultural (including family) environment (Koeppen-Schomerus et al, 2001). Yet, identifying the roles of different factors is important for the purposes of any policies aimed at dealing with the epidemic. If overweight is entirely genetic, then, short of a degree of genetic manipulation that is likely to be both technically infeasible and socially unacceptable, there is only a limited set of policy options available (Manski, 2012). If, on the other hand, there is a significant

¹ Public Health England *Child Weight Data Fact Sheet* August 1914.

http://www.noo.org.uk/securefiles/141007_1330//ChildWeight_Aug2014_v2.pdf

² Public Health England. http://www.noo.org.uk/NOO_about_obesity/child_obesity/UK_prevalence

cultural or environmental component in transmission, then there is room for policy intervention; but that component needs to be identified so that policy can be properly targeted.

Identifying the role of parents seems particularly important. It is possible that the spread of overweight among children can be attributed in large part to the influence of parental norms, including unhealthy role modelling. Children may consciously or unconsciously observe and model their parents especially with regards to fitness and to food consumption. Indeed, there is evidence that children's caloric intake, diet habits, level of physical activity and health behaviour in general are, at least partially, dictated by their parents' health behaviour and culturally determined social norms (Anderson and Butcher, 2006).

In this paper, we address the question concerning the existence and magnitude of the parental cultural influence on children's overweight in England. Our paper contributes to the existing literature by shedding some light to the question of how transmittable are overweight and obesity. Although Sacerdote (2007) found little evidence that that overweight is transmitted from parents to adopted children, the generalizability of these findings may be limited by the fact that the study uses a quasi-random design focusing on Korean adoptees in the US and that Koreans rates of obesity and overweight are among the lowest of the world (OECD, 2015). On the other hand, Koeppen-Schomerus et al. (2001) uses twin studies and provides evidence that overweight is not highly heritable. To speak to such debate, our study draws upon all the thirteen waves of the Health Survey for England (HSE) to construct a unique dataset containing children living in homes with either two biological parents or two adoptive parents. Besides the nature of the child-parent relationship, the data include information on a range of children's and parents' characteristics; on parental lifestyles; and on validated anthropometric records on children's overweight. These data allow us to identify the magnitude of the cultural transmission of overweight and obesity by quantifying the differences in the degree of transmission from parents to children between those

children living with two biological parents and those *living with two strictly³ adoptive parents*. Our estimates control not only for children's characteristics, parents' traits and other common environmental factors, but also for sample selection bias resulting from adoption not being a random event, with some sorts of households being more likely to adopt a child than others. Additionally, we contribute to a contentious point in the literature about whether maternal full-time employment alters the transmission of overweight, even when genetic transmission is not having an effect.

Our results reveal that when both adoptive parents are *overweight*, the likelihood of an adopted child being *overweight* is between 10% and 20% higher than when they are not, a result that we attribute to *cultural/environmental transmission* of overweight. We also find that the *cultural transmission*⁴ of *overweight* from parents to children is not aggravated by having a full-time working mother. Nevertheless, for natural children only, having a full-time working mother does significantly increase the positive effect of having an obese father on the likelihood of the child being overweight or obese.⁵

The remainder of the paper is organized as follows. Section 2 contains the model and outlines the empirical strategy. Section 3 describes our dataset. Section 4 reports our results. Section 5 discusses them, and Section 6 concludes.

2. Background and Empirical Strategy

Our empirical strategy is grounded on a health production function framework that allows the differentiation of genetic and environmental mechanisms in the intergenerational transmission of

³ We exclude those living with genetically related adoptive parents.

⁴ Throughout the text, we refer to cultural and environmental transmission indistinguishably.

⁵ Our measure of overweight includes obesity.

overweight. Health and non-health related traits of the parental environment influence some of the arguments in the child's production function creating links between the two generations as in Thompson (2014), who studies the intergenerational transmission of health using a CES production function model (see also Cunha et al., 2010; Cunha and Heckman 2007; and, Todd and Wolpin, 2003). In our case, we adapt the model of health vertical transmission by letting o_i indicate the overweight condition of the child i , and g_i and e_i the genetic and environmental factors influencing the weight of a child, respectively, so that $o_i = A[ag_i\gamma + (1-a)e_i\gamma]^{1/\gamma}$. The factor g_i reflects both genes and the genetic predisposition to be overweight or obese. The factor e_i contains non-genetic influences, including socio-economic and environmental factors such as: age; gender; education; socio-economic and employment status; and urban versus rural dwelling. The intergenerational transmission stems from the fact that parents and children share with different degrees the arguments in the factors g_i and e_i . As Thompson (2014) illustrates for health, in our setting when $\gamma = 1$ genes and environment have an additively separable influence on overweight status of the child and α and $1-\alpha$ represent the relative weight that e_i and g_i have, respectively, in the likelihood of a child being overweight.. When $\gamma \leq 1$, there is no separability between genes and environment and they interact in the production of health.

In our setting, we assume that being overweight has both genetic and environmental (or cultural) causes and that, as for other conditions, the specific interaction of genes and environmental factors will be crucial in determining whether a child is overweight. For instance, a predisposition of the parents to gain weight arguably may make them more aware of the nutritional content of food or of the need to do exercise, and this may translate in their children being exposed to healthier foods and more exercise, and ultimately less likely to be overweight.

As we explain below, we present estimates of different econometric specifications that compare the transmission of overweight across biological and adopted children. The results of the estimation for

non-biological children should remove the shared genetic components of transmission in e_i . But, as pointed by Thompson (2014), since we are not able to identify the gene-environment interactions, the resulting estimates for adoptees represent the average of α over the support of e . Moreover, since assignment to a given type of household (both biological parents; only one biological parent; and both adoptive parents) is not random, correcting for observable and unobservable sample biases will be crucial to identify non-genetic transmission of overweight. We correct for these biases to the extent that we can both by using a Heckman selection model and by propensity score matching design.

Our empirical strategy is to estimate a reduced form specification that draws upon the health production function above. We specify a linear model in which the latent overweight of a child is explained by non-genetic factors (age of the parents, their education and employment statuses, household's income, type of dwelling, and, being exposed to passive smoke); the child's own characteristics (age, gender, ethnic group); and, indicator variables taking value 1 if both parents being overweight; only the mother being overweight; or only the father being overweight, respectively:

$$o_{ij}^* = \delta_0 + \delta_b o_{ij}^b + \delta_M o_{ij}^M + \delta_F o_{ij}^F + \beta Z_j + \theta X_{ij} + v_{ij} \quad , \quad (1)$$

where o_{ij}^* indicates the latent overweight of child i in household j ; o_{ij}^b is an indicator variable for *both parents* of child i in household j being overweight or obese; o_{ij}^M takes value one if *only the mother* of child i in household j is overweight ; o_{ij}^F takes value one if *only the father* of child i in household j is overweight ; Z_j is a vector with the parents' characteristics and X_{ij} a vector of the child's characteristics; and v_{ij} is the error term.⁶ Assuming normality of the error term, v_{ij} , the probability of

⁶ Note that, given the data available in HSE, for children living with their natural parents, o_{ij}^M and o_{ij}^F refer to the overweight status of their biological parents. For children living with their adoptive parents, these terms will refer to the overweight condition of the *adoptive* parents.

observing that a child i in our sample is overweight ($o_{ij} = 1$) is the probability that the corresponding latent variable is positive, i.e.:

$$P(o_{ij} = 1) = P(o_{ij}^* > 0) = \Phi(\delta_0 + \delta_b o_{ij}^b + \delta_M o_{ij}^M + \delta_F o_{ij}^F + \beta Z_j + \theta X_{ij}) \quad (2)$$

Therefore, in this framework, coefficients δ_b , δ_M , and δ_F will estimate the effect of *both parents, only the mother* or *only the father* being overweight on the likelihood a child being overweight, respectively.

We estimate equation (2) for two different groups of children: those who live with both biological parents and those who live with both adoptive parents. The difference between the coefficients for children that are biological (exposed to both genetic and environmental transmission of overweight and those that are adopted (only to the environmental transmission), will give us a measure of the relative importance of environmental intergenerational transmission for overweight.

We first estimate equation (2) using a Probit model, without taking into account the selection bias of children into each of these groups. Second, we perform robustness checks re-estimating equation (2) controlling for the sample selection bias of being in an adoptive family by using both a probit models with sample selection⁷ (heckprobit) and a propensity score matching-based correction. The *exclusion restriction* for the identification of the Heckprobit models relies on the parents' age and the father being unemployed, which are likely to affect the likelihood of an individual being adopted but not the overweight of the child. The propensity matching score corrects the effects of sample selection (as in Rosenbaum and Rubin, 1983) by allowing estimating the conditional probability of each child being in an adoptive household given observed covariates of the child and the household. The propensity score is then used as a covariate to adjust the original model. As an additional robustness check, we also estimate equation a variation of equation (2) using Ordinary Least Squares.

⁷ i.e., Maximum-Likelihood probit models with sample selection as in Van de Ven and Van Praag (1981).

Additionally, we estimate equation (2) allowing the mother working full time to influence the degree of transmission of overweight from parents to children. We do so by interacting the indicator variable taking value 1 when the mother works full time with the overweight indicator variables for the parents.⁸

We have considered additional specifications including the specific transmission of mother-daughter and father-son; and whether the transmission has evolved with each wave of the survey, i.e. over time. We do not include these results as sample limitations hampered the robustness of the coefficients.

3. Data

The dataset we use to estimate the models above originates in the Health Survey for England (HSE). The HSE is an annual cross-sectional survey designed to measure health and health-related behaviours, including weight and height, body mass index (BMI), fruit and vegetable consumption, alcohol consumption and smoking in adults and children living in private households in England. The survey also contains the socio-economic status of the household and core information on all its members, including their relationship. This allows us to categorize children in types of households depending on whether they live with both their biological parents or they live with a set of parents neither of whom is biological and unrelated genetically.⁹ Our pooled cross-section panel dataset results from merging information contained in thirteen different waves of the HSE, from 1997 to 2009.

Adoption in the UK can be legally carried out by parents that are over 21 years of age that have at least one year of residency and have a fixed permanent home in the UK irrespective of the civil status. The latter includes the possibility of the partner of the natural parent to being considered

⁸ We also estimated the model using families in which one of the parents is biological and the other is not but given that the baseline characteristics of this type of households are markedly significantly different from the natural and adoptive parents' families, we do not present it in here.

⁹ As we have the relationship between children and all relatives in the household, our sample does not include children living with 'non-parents' but biologically related family members, i.e. grandparents, uncles, aunts, etc.

‘adopter parent’ too (UK Government, 2013)¹⁰. The process take place after an application to an adoption agency whether a council or a privately run one. The conditions to be met to be regarded as suitable include a *full medical examination*, a police check of no pre-existing convictions, including three-reference letters, training and an assessment by a social worker. Recommendations regarding suitability of an adopter parent are made by an external ‘adoption panel’. Once an adoption panel makes decisions, then the parents are matched with a child locally or referred to the Adoption Registry.

Because of the nature of our dataset, we are confronted with several limitations. First, we do not have information on the biological parents of the adopted children. Thus, we cannot control for early nutrition effects they may have faced and we cannot observe the weight of the biological parents. Second, we cannot identify the exact time of adoption, and can only indirectly control for it through age. Third, we cannot identify whether if the individuals were born overseas although we do have their ethnicity information.

More generally, studies using data from adoptees face challenges that complicate the identification strategy (Holmlund et al, 2011). Parental sorting is not random. “[A]doption agencies often place infants selectively by matching natural and adoptive parent characteristics, such as education, occupation, and impressions about intelligence” (Scarr and Weinberg, 1994). Thus, if the genetic influence of the biological parents is not accounted for, statistical associations between the outcomes of adopted children and their adoptive parents could reflect a combination of the adoptive parents’ environmental influences and the correlated genetic inheritance. A way to partially address this is to correct for sample selection into adoptive families using the characteristics of the child and the foster parents. Using this approach, Bjorklund *et al* (2006) find no evidence of the existence of a sample selection bias as estimates between adoptees and biological parents in Sweden; Sacerdote (2007) uses

¹⁰ <https://www.gov.uk/child-adoption>

a sample of American Korean adoptees quasi-randomly assigned to adoptive families and finds evidence of cultural transmission of some health behaviours and BMI. In our case, we go extend the analysis in several directions: First, we use a measure of overweight and obesity of both parents and children obtained by a nurse during the survey instead of relying on BMI. Having socioeconomic information of both parents for all children allows to control for the potential compounding effect of assortative mating. Second, we are able to correct for potential sample selection biases based on observables due to selective adoptee placement and the different characteristics of the adoptive families. Third, we run a battery of subsample analyses and robustness checks to investigate the stability of our estimates.

In this paper, we limit the source of disparity between our sample of biological and adoptive families by restricting our analysis to two-parent households. We also use children's and parents' observable characteristics to correct for sample selection biases exploiting Heckprobit models¹¹ and Propensity Score Matching (PSM) based ones.

Our final dataset contains children of all waves, including their socio-demographic characteristics, their physical measurements (BMI, weight, height, etc.), those of their parents and the nature of their relationship. The measurements of height and weight in the HSE are validated by a nurse, thus overcoming the problem of measurement error of these values present in other surveys containing children, i.e. Phipps et al. (2004) or Anderson et al. (2004).

[Insert Table 1 about here]

Table 1 provides our sample descriptive statistics including the rates of overweight and obesity for children and their parents. We report the statistics for the overall sample (13,836 observations), and

¹¹ as in Van de Ven and Van Praag (1981).

disaggregated by type of household, i.e. those in which both parents are biological (13,536 observations) and those in which both parents are adoptive (300 observations). In the last column we show the outcome of the T-Tests analysing if the means of the two groups are significantly different.

Looking at these statistics and the results of the T-tests, we observe that only for nine out of forty-eight variables is the difference between the groups statistically different at the 99% level and for five variables the difference is significant at the 90% level. In the light of this, we are confident that the baseline characteristics of our biological and adopted household are not challengingly different. We do observe nevertheless that adopted children in the sample are slightly older than those in a biological parents' household; they are slightly more likely to have an obese mother, an obese father, or both parents obese; their parents tend to answer the education question less often and when they do, they are less likely to be in the lower end of the education distribution. Their mothers choose the 'other' occupation category more often; their parents are slightly older; they live less often in suburban areas; and, they are more often exposed to passive smoking.

The percentage of overweight children is about 23% (slightly higher for adopted but not statistically significant); of obese children 5.6%; of both parents being obese, 7% for the biological parents' households and 10% for the adoptive; of both being overweight ,about 40% for the former type of household and 47% for the latter. Only the mother being obese happens in about 16% of our sample; only the father being obese in 15% of the first type of households and in 17% of the second type (but again the difference is not statistically significant); only the mother being overweight in about 13% of the biological parents' families and in 11% of the adoptive families. Lastly, only the father is overweight in about 30% of both types of households. These univariate differences in the percentage of obese and overweight parents could be due to the slightly higher age of adoptive parents. We refer to the table for further details on the exact figures for the forty-eight variables. Finally, it should be noted that unlike BMI in adults, BMI among children changes over time and hence fixed thresholds

can provide misleading findings. Hence, for the children we use the international standard BMI cut off points for age and sex published by the International Obesity Task Force (IOTF) as in Saxena et al. (2004). For parents, we used the standard overweight and obesity BMI cut-offs: parents are classified as overweight if their BMI is between 25 and 30 and as obese if it is greater than 30.

4. Results

Results are presented in Tables 2, 3 and 4. Table 2 shows the estimates of the transmission of the both parents being overweight on the likelihood of the child being overweight. The dependent variable is indicated in the top row and whether the parents are overweight is indicated in the second row. The third row in this table indicates which type of household the child is living in (both parents biological or both parents adoptive). The method used to estimate these coefficients is a probit model and expressed as marginal effects. Table 3 re-estimates the coefficients in Table 2 by correcting the sample selection potential biases of belonging to each type of household using Heckprobit models and propensity score matching models. Finally, Table 4 is an extension of all preceding tables in which the effect of the parents' weight on that of their children is estimated controlling for the fact that the mother works full time.

[Insert Table 2 here]

The results in the first two columns of Table 2 indicate that the transmission *overweight from parents to children* is significant and positive when both parents are overweight for both groups of families. The increase of the likelihood of being overweight of those children when both parents are biological is 0.270,¹² and for those adopted 0.210. Given that the biological-parents coefficient is picking up both genetic and cultural transmission, whereas the adopted-parents coefficient only reflects cultural

¹² Throughout the paper, results express percentage points, i.e. 0.116 means an increase of 11.6 percentage points and *an increase of 12%* refers to an increase in 12 percentage points.

transmission, this suggests that the relative importance of the cultural transmission when both parents are overweight is big. *Only the mother being overweight* increases significantly the likelihood of the offspring being overweight by 12% only for children living with both biological parents, but not for the adopted group. *Only the father being overweight* is significant both for families where both parents are biological (0.116) and for those where they are adoptive (0.240). The difference between these two coefficients suggests that, when only the father is overweight, the *cultural transmission* for adopted children is more important than both the genetic and cultural transmission for natural children.

In the second panel we report the estimates of the effect of the *parents being obese* on the probability of the *children being overweight*. For those with both biological parents, *both parents being obese* increases the likelihood of the children being overweight by 0.342; *only the mother being obese* by 0.176; and *only the father being obese* increases it by 0.144. For those families in which both parents are adoptive, the only significant coefficient is that of *both parents being obese* and its effect on the probability of the child being overweight is 0.216. The cultural intergenerational transmission of obese parents to overweight children thus seems a bit weaker than of overweight parents to overweight children, but still very sizeable.

Finally, the third panel in Table 2 looks at the relationship between the *obesity of the parents* on the probability of the *children being obese*. For this case, for the first type of families (both biological), if *both natural parents are obese*, the likelihood of the *child being obese* as well increases by 0.170, when *only the mother is obese*, it increases by 0.070 and when *only the father is obese* by 0.044. For the adoptive families, if *both parents are obese*, the likelihood of the child being obese goes up by 0.208 but the effects of the *only the mother* or *only the father* being obese are not significant probably due to the small sample size. So again there appears to be cultural transmission of obesity, but by a smaller proportion than overweight.

[Insert Table 3 here]

Table 3 corrects the estimates in Table 2 by sample selection using two-stage Heckprobit and PSM-based models.¹³ Results in Table 3 are quite similar to those in Table 2 but a few remarks are to be made:

First, in the Table 3, the estimates of the effects of *both parents being overweight* on the likelihood of the *child being overweight* are higher than when not correcting for sample selection for those households where both parents are adoptive (above 0.246 instead of 0.210 in Table 2), and slightly smaller for those living with their biological parents (0.252 instead of 0.270). The effect of *only the mother being overweight* and *only the father being overweight* become significant for adopted children when using the PSM-based correction for sample selection (0.154 and 0.106, respectively); using the Heckprobit correction *only the father being overweight* is significant and higher than in Table 2 (0.272 instead of 0.240). The bias due to sample selection of the adoptive households does not appear to be large for the overweight estimates judging by the similar estimates in the first panel in Table 3.

Second, in the second panel corresponding to the influence of *obese parents* on the *child being overweight*, we observe that the sample selection correction increases all coefficients for the biological parents' households and also for adoptive children if the PSM-based approach is applied, but not for the Heckprobit correction approach. Also, the PSM-based estimates coefficients of *only the mother* or *only the father being obese* for adoptive children become significant and similar to

¹³ The selection equation in the Heckprobit model includes parental age and the father being unemployed. The PSM scores use gender of the child, whether the mother works full time, the father being unemployed, household size, and whether the household lives in an urban setting.

those of the biological parents households (0.166 and 0.139, respectively). The Heckprobit model estimates of these indicator variables remain insignificant.¹⁴

Third, by looking at the third panel, we observe that sample selection correction by either approach reduces slightly the effect of the *transmission of obesity from parents to children* for children living with their biological parents. Using the Heckprobit correction model reduces greatly the effect of both adoptive parents being obese on the probability of the adoptee being obese (0.027). The sample selection correction using the PSM-approach increases the effect of *both parents being obese* (to 0.181) and *only the mother being obese* or *only the father being obese* are significant and positive (0.082 and 0.051, respectively).

Finally, the fact that the PSM-scores interacted with our variables of interest are not significantly different than zero could indicate that the sample selection issues in our sample are not excessively worrying, which is in accordance with the baseline characteristics of our two samples being quite similar. Thus, the biases in the coefficients reported in Table 3 should not be too important.

[Insert Table 4 here]

The results in Table 4 also test whether the fact that the mother works full time has an impact on the overweight transmission estimates. To do so, we estimate the specifications in Table 2 allowing for an interaction of an indicator variable of the mother working full time with the overweight/obesity status of the parents. As can be observed from the table, none of the interactions are significant except for that with the *obesity status of the father only* in the second and third panel and only for the biological parents' kind of families. Thus, when *only the father* is obese and at the same time the *mother works full time* the likelihoods of both the child being overweight (0.047) and being obese (0.020) increases significantly beyond the sole effect of the father being obese. But, probably due to

¹⁴ The fact that the PSM correction based estimates are more similar to the natural parents' estimates in panel 2 and 3 could be explained by the fact that the PSM corrects for fewer observables than the heckprobit.

sample size issues arising from the interaction terms, some of the coefficients that were significant in previous specifications for adopted children are insignificant when using this specification.

5. Discussion

Overweight is an expression of both genetic and cultural influences. In this paper we have attempted to estimate the cultural transmission of overweight. We contribute to the literature of intergenerational transmission of health, by quantifying the strength of the intergenerational correlation of overweight in both natural children and adoptees. The analysis is conducted making use of a uniquely constructed dataset of English adoptees from 1997 to 2010. We have examined intergenerational transmission alongside a long list of other confounding variables that could be driving the association such as education, parental and child age, gender effect and, following the literature, the effect of female labour market participation.

We base our empirical approach on a theoretical model of health production by which children's overweight depends on the overweight or obese status of their parents, and thus implicitly on the parents' lifestyle choices and net caloric intakes. We follow an empirical strategy that has taken selection issues in consideration alongside drawing upon a naïve probit model. We estimate our empirical models of overweight for two types of children, those living with both their natural parents and those living with adoptive parents. We use various specifications, which include the observable characteristics of the child and the parents.

Our results indicate quite strongly that there seems to be a powerful cultural transmission of overweight inter-generationally, in addition to that resulting from the genetic links even when we control for sample selection employing two different strategies using observables. For obesity, the results are less strong, but both parents being obese or the father alone being obese, increase the

probability of observing an overweight and/or an obese child even when they are not genetically related. However, the mother alone being obese is an insignificant factor.

These findings are robust to different specifications, including the mother working full time and income, which has been pointed out as the culprit for child's obesity (Anderson, 2003). We do not find evidence that the mother works full time explains children's obesity, nor their tendency to be overweight. We control for education of both parents, type of dwelling, various characteristics of the household, and degree of urbanisation. Our findings survive the inclusion and exclusion of these controls.

There is an intriguing aspect to these results. In general, the results concerning the powerful cultural transmission effect are much stronger for overweight than obesity. If both adoptive parents are overweight, or if only the father is overweight, this increases the probability of the children being overweight by about 25% to 30%. However, if both adoptive parents are obese, when we control for the mother working full time this has no significant effect on children's obesity. This suggests that the primary mechanism of the intergenerational transmission of obesity is much more likely to be genetic than that for overweight. Indeed, we can find little evidence from our results of any important cultural transmission of obesity.

The importance of the cultural transmission of overweight may be emphasized by the fact that in some of the specifications, when correcting the sample selection bias of adoptive children using the PSM approach, *both adoptive parents being overweight* has a larger impact on the probability of their children being overweight than when *both biological parents are overweight*. This would suggest that natural parents would have a far smaller cultural impact on their children being overweight than adoptive parents do. The latter can be the result of their being more likely to follow a different lifestyle pathway unrelated to biological triggers of behaviour.

Another thought-provoking feature of the results concerns a difference in the impact of the non-natural mother's and father's overweight. In some of the specifications the mother's overweight is not significant while the father's is. A possible explanation is that the mother is in charge of the nutrition of the children and their father and may tend to overfeed them while underfeeding or feeding adequately herself.

6. Conclusion

This paper has drawn upon a uniquely constructed dataset of English adoptees to investigate the existence and mechanisms of intergeneration transmission of overweight. We have found that that children's overweight is robustly related to the overweight of the parents, even when there is not genetic transmission as is the case of adoptees. However, while we can establish there is a strong cultural transmission of overweight, our evidence is weaker for obesity.

We also find that the *cultural transmission of overweight* or *obesity* from parents to children is not aggravated by having a full-time working mother. Nevertheless, for natural children only, having a full-time working mother does significantly increase the positive effect of having an obese father on the likelihood of the child being overweight or obese.

We acknowledge that our estimates are subject to several limitations imposed by the nature of the data. First, adopted children might belong to a healthier/unhealthier sample than the biological, although a wealth of studies suggest that selective placement of adoptees does not seem to have an impact on the cultural transmission of health (Wilcox-Gok, 1983) and thus on health itself. Second, although adopted children are not genetically related to their parents, adoption agencies do attempt to

match biological and adoptive parents in various ways (selective placements), a factor that could cause additional sources of sample selection.¹⁵ Third, we cannot observe the age of adoption (though the majority of adoptions takes place before the age of 3) and, hence, we cannot control for the length of a child's exposure to his/her adoptive family environment. Fourth, unlike the data obtained from adoption registers, we do not have information on the biological parents of the adoptees, and whether the children were foreign born or not. To address some of the non-randomness issues, we have compared the two types of households to ensure they are not significantly too different and still correct for sample selection biases using two-stage Heckman models and Propensity Score Matching (PSM) adjustments. We have also run robustness checks using different specifications. Finally, the sample of obese adopted children is small, and the number of those who have obese parents even smaller. This hinders the strength of our results regarding the cultural transmission of obesity from parents to children.

Our paper improves upon existing literature by using the Health Survey for England to examine a sample of children living in homes where parents are either both adoptive or both biological. The advantage of this dataset is that it contains the same data on adopted and biological children and their living-in parents, including anthropometric measurements and parents', children's and household's characteristics. Thus, unlike data on adoptees from administrative records, we do not need to match the sample of children with the general population.

A comparison of our findings with that of the wider literature on intergeneration transmission for education (Holmlund et al, 2011) reveals that for obesity genes play a larger role than for overweight, which is quite sensitive to changes in the environment. This is consistent with health conditions such as asthma, allergies, headaches and diabetes (Thomson, 2014) and other studies that do not

¹⁵ To address this issue some studies use information on the adoptees' biological parents (Björklund et al, 2006), and Sacerdote (2007) draws upon a random assignment of children.

disentangle total from cultural transmission (Classen and Hokayem 2005, Classen, 2010 and Costa-Font and Gil, 2013).

We conclude that this paper provides evidence in favour of the hypothesis that there is a strong cultural component in the transmission of cultural habits that promote overweight from parents to children. That is, gender specific effects might still reflect that, as some studies show (Lake et al., 2006), food responsibility was predominately a female dominated, but the ingest of such food might be more than proportionally consumed by men and children. The importance of *both* parents being overweight in explaining the overweight of the children might as well reflect evidence of assortative mating, or alternatively a reinforcing environmental effect that takes place when both parents adopt similar behaviours. One hypothesis consistent with assortative mating is that health and lifestyle preferences end up determining partner-matching. Thus, both parents may be overweight or obese as a result of sharing a common lifestyle and tastes, which are in turn passed on to their children.

Our results suggest that there is room to design policies to tackle children's overweight and obesity by influencing parental overweight and their lifestyles, and that ideally both parents should be influenced for the effect to be more effective; otherwise problems of children overweight are likely to persist. Overweight is passed through generations, and the pathway seems to be primarily driven by the children environment. In contrast, and consistently with the behavioural genetics literature, obesity exhibits a highly genetic component.

References

Ahlburg, D., 1998. Intergenerational transmission of health. *The American Economic Review Papers and Proceedings* 88 (2), 265–270.

Anderson, P, Butcher, K and Levine, P (2003). Maternal Employment and Overweight Children. *Journal of Health Economics*, 22(3): 477-504.

Anderson, P., Butcher, K., Schanzenbach, D. (2007). Childhood disadvantage and obesity: is nature trumping nurture? *NBER Working Paper No. 13479*.

Anderson, PM and Butcher, KF (2006). Childhood obesity: trends and potential causes. *The Future of Children*, 16(1): 19-45.

Amelia A. Lake, Robert M. Hyland, John C. Mathers, Andrew J. Rugg-Gunn, Charlotte E. Wood, Ashley J. Adamson, (2006) "Food shopping and preparation among the 30-somethings: whose job is it? (The ASH30 study)", *British Food Journal*, Vol. 108 Iss: 6, pp.475 – 486

Bisin, A. and Verdier, T. (2001), 'The Economics of Cultural Transmission and the Dynamics of Preferences', *Journal of Economic Theory* 97(2), 298-319.

Bisin, A and Topa, G (2003). Empirical Models of Cultural Transmission, *Journal of the European Economic Association*, 1(2-3): 263-375.

Bisin, A., and T. Verdier (2000), “Beyond the Melting Pot: Cultural Transmission, Marriage, and the Evolution of Ethnic and Religious Traits,” *Quarterly Journal of Economics* 115(3), 955-988

Björklund, A,M Lindahl, and EPlug (2006). “The Origins of Intergenerational Associations: Lessons from Swedish Adoption Data.” *Quarterly Journal of Economics* 121:999-1028.

Björklund, Anders, and Laura Chadwick(2003). “Intergenerational Income Mobility in Female Labor Force Dynamics,” *Quarterly Journal of Economics*, 119(4), 1249-1299.

Björklund, Anders, Markus Jäntti, and Gary Solon (2005). Influences of Nature and Nurture on Earnings Variation: A Report on a Study of Various Sibling Types in Sweden. In Samuel Bowles, Herbert Gintis, and Melissa Osborne Groves,(eds) *Unequal Chances: Family Background and Economic Success*. Princeton: Princeton University Press, 2005, pp. 145-164.

Case A, Fertig A, Paxson C (2003) ‘*From Cradle to Grave? The Lasting Impact of Childhood Health and Circumstance*’ NBER Working Paper 9788

Cesarini, D, Dawes, C.T ,Johanesson, M Lichtenstein, P and Wallace, B (2009a). Genetic Variation in Preferences for Giving and Risk Taking. *Quarterly Journal of Economics*. *Quarterly Journal of Economics*, 124: 809-842

Cesarini, D., Lichtenstein, P., Johannesson, M. and Wallace, B. (2009b), Heritability Of Overconfidence. *Journal of the European Economic Association*, 7: 617–627

Chia, Yee Fei. 2008. “Maternal labour supply and childhood obesity in Canada: evidence from the NLSCY” *Canadian Journal of Economics*, 41(1):217-242.

Classen, T., Hokayem, C., (2005). Childhood influences on youth obesity. *Economics and Human Biology* 3(2), 165–187.

Classen, Y (2010). Measures of the intergenerational transmission of body mass index between mothers and their children in the United States, 1981–2004. *Economics and Human Biology* 8 30-43.

Coate, D (1983). *The relationship between diet, parent's fatness and obesity in children and adolescence*. NBER Working Paper, 1072.

Coate, D. (1983) *The relationship between diet, parent's fatness, and obesity in children and adolescents*, NBER Working Paper No, 1072.

Coneus, K and Spiess, CK (2011). Intergenerational Transmission of Health in Early Childhood – Evidence from the German Socio-economic Panel Study. *Economics and Human Biology*, 10: 89-97.

Costa-Font, Joan and Gil, Joan (2013) *Intergenerational and socioeconomic gradients of child obesity* *Social Science and Medicine*, 93. 29-37.

Currie J and Moretti E (2007) 'Biology as Destiny? Short- and Long-Run Determinants of Intergenerational Transmission of Birth Weight' *Journal of Labour Economics*, Vol 25 No. 2, pp 231–264.

DeAgostini, P (2007). *Diet Composition, Socio-Economic Status and Food Outlets Development in Britain*. ISER Working Paper 2007-2009.

Duncan, Haller and Portes (1968) Peer influences on aspirations: A Reinterpretation. *American Journal of Sociology* 74(2):119-37.

Duncan, G, A Kalil, S E. Mayer, R Tepper, and M R. Payne (2005). The Apple Does Not Fall Far from the Tree. In *Unequal Chances: Family Background and Economic Success* Edited by Samuel Bowles, Herbert Gintis, & Melissa Osborne Groves, Princeton University Press.

Elks CE, den HM, Zhao JH, Sharp SJ, Wareham NJ, Loos RJ, et al (2012). Variability in the heritability of body mass index: a systematic review and meta-regression. *Front Endocrinol*, 3:29

Emanuel I, Filakti H, Alberman E, Evans SJ.(1992) Intergenerational studies of human birthweight from the 1958 birth cohort. 1. Evidence for a multigenerational effect. *Br J Obstet Gynaecol* 1992;99:67–74.

Fernández, R., A. Fogli, and C. Olivetti (2004), “Mothers and Sons: Preference Formation and Permanent and Separated Families.” *Economics Letters* 80, 239-246.

Fertig, Angela, Gerhard Glomm, and Rusty Tchernis (2009). “The connection between maternal employment and childhood obesity: inspecting the mechanisms” *Review of Economics of the Household*, 7:227-255

Fogelholm M, Kukkonen-Harjula K (2000). Does physical activity prevent weight gain : a systematic review. *Obes Rev* 2000; 1:95–111.

Goode, A., Mavromaras, K. G. and Smith, M. (2008). *Intergenerational Transmission of Healthy Eating Behaviour and The Role of Household Income*, IZA Discussion Paper No. 3535

Goran MI, Shewchuk R, Gower BA, Nagy TR, Carpenter WH, Johnson RK (1998). Longitudinal changes in fatness in white children effect of childhood energy expenditure. *Am J Clin Nutr* 1998; 67: 309–316.

Greenough, W.T, Black, JE and Wallace, CS (1987). Experience and brain development, *Child Development*, 58: 539-59.

Grilo CM, Pogue-Geile ME. (1991). The Nature of Environmental Influences on Weight and Obesity: A Behaviour Genetic Analysis. *Psychological Bulletin* 110(3); 520-537.

Hamermesh, D (2010). Incentives, time use and BMI: The roles of eating, grazing and goods. *Economics and Human Biology* 8 , 2–15.

Hebebrand J., Wulftange H., Goerg T., Ziegler A., Hinney A., Barth N., Mayer H., Remschmidt H. (2000). Epidemic obesity: are genetic factors involved via increased rates of assortative mating? *Int. J. Obes. Relat. Metab. Disord.* 24, 345–353

Hruschka, D.J and AA. Brewis (2013). Intercontinental differences in overweight of adopted Koreans in the United States and Europe', *Economics and Human Biology* 11 (3), 2013, pp. 345-350.

Holmlund, H, Lundhal, M and Plug, E (2011). The Causal Effect of Parents Schooling on Children's Schooling : A Comparison of Estimation Models. *Journal of Economic Literature*, 49 (3): 615-651.

Koeppen-Schomerus, G J Wardle and R Plomin (2001). A genetic analysis of weight and overweight in 4-year-old twin pairs *International Journal of Obesity* (2001) **25**, 838-844

Llewellyn CH, Trzaskowski M, Plomin R, Wardle J (2013). Finding the missing heritability in pediatric obesity: the contribution of genome-wide complex trait analysis. *Int J Obes* 37(11):1506-9.

Liu, E., Hsiao, C., Matsumoto, T., Chou, S (2009) , Maternal full-time employment and overweight children: Parametric, semi-parametric, and non-parametric assessment. *Journal of Econometrics*.

Maes HHM, Neale MC, Eaves LJ. (1997) Genetic and Environmental Factors in Relative Body Weight and Human Adiposity. *Behavior Genetics* 27(4); 325-351.

Manski, C (2011). Genes, Eyeglasses, and Social Policy. *Journal of Economic Perspectives*, 25 (4): 83- 9.

Martin, M., (2008). The intergenerational correlation in weight: how genetic resemblance reveals the social role of families. *American Journal of Sociology* 114, S67–S105

Martin Caraher, Paul Dixon, Tim Lang, Roy Carr-Hill, (1999) "The state of cooking in England: the relationship of cooking skills to food choice", *British Food Journal*, Vol. 101 Iss: 8, pp.590 - 609

Moorman, J.E., and Hernandez, D.J (1089). Married-couple families with step, adopted, and biological children. *Demography* (May 1989) 26(2):267-77

OECD (2015). Obesity and the Economics of Prevention: Fit not Fat - Korea Key Facts. Organisation of Economic Cooperation and Development (<http://www.oecd.org/els/health-systems/obesityandtheconomicsofpreventionfitnotfat-koreakeyfacts.htm>, accessed January 2015).

Oreopolous P, Stabile M, Walld R, Roos L 'Short, Medium, and Long Term Consequences of Poor Infant Health: An Analysis using Siblings and Twins' NBER Working Paper No. W11998.

Parsons, TJ. Parsons1, C Power1 and O Manor (2005). Physical activity, television viewing and body mass index: a cross-sectional analysis from childhood to adulthood in the 1958 British cohort. *International Journal of Obesity* (2005) 29, 1212–1221.

Phipps, Shelly A., Peter Burton , Lynn Lethbridge, and Lars Osberg. 2004. “Measuring Obesity in Young Children.” *Canadian Public Policy*. 30(4):349-364.

Plomin R. (1990). The role of inheritance in behavior. *Science* 248, 183–188

Plug, E (2004). Estimating the Effect of Mother’s Schooling on Children’s Schooling Using a Sample of Adoptees.” *American Economic Review* 940 , 358-368.

Plug, Erik. and W Vijverberg (2005). “Does Family Income Matter for Schooling Outcomes? Using Adoptees as a Natural Experiment.” *Economic Journal* 115, 879-906

Propper C., Rigg J., Burgess S. (2007) Child health: evidence on the roles of family income and maternal mental health from a UK birth cohort. *Health Economics* 16(11): 1245-1269

Rosenbaum P.R., Rubin D.B., (1983) The Central Role of Propensity Score in Observational Studies for Casual Effects, *Biometrika*, 70, 41–55

Rhee K.E., Phelan S., McCaffery J, (2012) ‘Early Determinants of Obesity: Genetic, Epigenetic, and In Utero Influences’, *International Journal of Pediatrics*, Vol 2012.

Sacerdote, B (2007). "How Large Are the Effects from Changes in Family Environment? A Study of Korean American Adoptees. *Quarterly Journal of Economics* 122, 119-157.

Saxena S., G Ambler, T J Cole, A (2004). Majeed 'Ethnic group differences in overweight and obese children and young people in England': cross sectional survey , *Arch Dis Child* 2004;89:30-36

Scarr, Sandra, and Richard A. Weinberg. "Educational and Occupational Achievements of Brothers and Sisters in Adoptive and Biologically Related Families." *Behaviour Genetics* 24 (1994), pp. 301-325.

Silventoinen K., Kaprio J., Lahelma E., Viken R. J., Rose R. J. (2003). Assortative mating by body height and BMI: Finnish twins and their spouses. *Am. J. Hum. Biol.* 15, 620-627.

Sorensen, T, I.A , Price, RA, Stunkard, AJ and Schulsinger, F (1989). Genetics of obesity in adult adopted and their biological siblings. *BMJ*, 298, 87-90.

Stunkardt, AJ, Thorkild I.A. Sørensen, Dr.med., Craig Hanis, Ph.D., Thomas W. Teasdale, M.A., Ranajit Chakraborty, Ph.D., William J. Schull, Ph.D., and Fini Schulsinger (1986). An adoption study of human obesity, *NEJM*, 314, 193-8.

Stunkard, AJ J R. Harris, N L. Pedersen, and GE. McClearn(1990). The Body-Mass Index of Twins Who Have Been Reared Apart. *N Engl J Med* 1990; 322:1483-1487

Townsend N (2009) *Obesity and Overweight Surveillance in England: what is measured and where are the gaps?* Oxford: National Obesity Observatory, 2009

Thomson, O (2014). Genetic mechanisms in the intergenerational transmission of health. *Journal of Health Economics*, 35: 132-145.

Van de Ven, W.P. and Van Praag, B.M. (1981). The demand of deductibles in private health insurance: A probit model with sample selection, *Journal of Econometrics*, 17, 229-252.

Von Hinke Kessler Scholder, S (2007). *Overweight Children: Does Timing Matter?* CMPO working Paper, 07/180.

Wardle J., Carnell S., Haworth C.M.A, Farooqi I.S., O'Rahilly S., and Plomin R. (2008), “Obesity associated genetic variation in FTO is associated with diminished satiety,” *Journal of Clinical Endocrinology and Metabolism*, vol. 93, no. 9, pp. 3640–3643.

Whitaker, R., Wright, J., Pepe, M., Seidel, K., Dietz, W., (1997). Predicting obesity in young adulthood from childhood and parental obesity. *The New England Journal of Medicine* 337 (13), 869–873

WHO (1999). Definition, diagnosis and classification of diabetes mellitus and its complications: Report of the WHO consultation. Geneva, World Health Organization (WHO), 1999.

Wickrama, KAS Rand D. Conger, Lora Ebert Wallace, Glen H. Elder, Jr (1999). The Intergenerational Transmission of Health-Risk Behaviours: Adolescent Lifestyles and Gender Moderating Effects. *Journal of Health and Social Behaviour*, 40(3): 258-272.

Wilcox-Gok, V (1983). The Determination of Child Health: An Application of Sibling and Adoption Data. *Review of Economics and Statistics*, 65(2): 266-73

Table 1: Summary Statistics and test of differences in means

		A	B		
		Overall Sample	Natural	Adopted	Sign Diff B-A
		Number of Observations		13836	13536
Child overweight and other characteristics	Obese Child	5.7%	5.7%	5.7%	
	Overweight Child	23.5%	23.5%	25.3%	
	Age of Child	9.1	9.0	10.9	***
	Female	49.1%	49.2%	46.3%	
	White	78.7%	78.6%	79.3%	
	Black (Caribbean, African or Other)	4.4%	4.4%	5.0%	
	South East Asian and Other Pakistan/Bangladesh/Chinese	12.7%	12.7%	15.0%	
		4.2%	4.3%	0.7%	
Parents' obesity	Obese Mother	21.1%	21.0%	26.0%	*
	Obese Father	22.4%	22.3%	27.3%	*
	Overweight Mother	13.0%	13.0%	11.3%	
	Overweight Father	31.7%	31.7%	29.3%	
	Both parents Obese	7.0%	6.9%	10.0%	*
	Only Mother Obese	14.1%	14.1%	16.0%	
	Only Father Obese	15.4%	15.4%	17.3%	
	Both parents Overweight	39.9%	39.7%	46.7%	*
	Only Mother overweight	13.0%	13.0%	11.3%	
	Only Father Overweight	31.7%	31.7%	29.3%	
Parents' characteristics	Mum Education: NA	13.1%	13.1%	15.3%	
	Mum Education: HE	31.2%	31.2%	32.3%	
	Mum Education: A/O Level	47.8%	47.8%	45.7%	
	Mum Education: CSE	5.9%	5.9%	5.3%	
	Mum Education Foreign	2.0%	2.0%	1.3%	
	Dad Education: NA	15.0%	14.8%	20.3%	***
	Dad Education: HE	41.3%	41.5%	32.7%	***
	Dad Education: A/O Level	37.2%	37.1%	39.0%	
	Dad Education: CSE	5.4%	5.4%	5.7%	
	Dad Education Foreign	1.1%	1.1%	2.3%	
	Mother at home	26.1%	26.2%	23.7%	
	Mother Employed	69.8%	69.8%	68.3%	
	Mother Retired	0.1%	0.1%	0.0%	
	Mother Other	4.1%	4.0%	8.0%	***
	Dad at home	1.3%	1.3%	2.0%	
	Dad Employed	90.4%	90.4%	88.3%	
	Dad Retired	0.7%	0.7%	1.7%	
	Dad Other	7.6%	7.6%	8.0%	
	Mother's Age	38.3	38.3	41.1	***
	Father's Age	41.0	40.9	43.8	***
Other Household	Income	£30,899.11	£30,913.34	£30,257.37	

characteristics	Own Flat	82.7%	82.7%	84.0%	
	Small Family	44%	45%	13%	
	Large Family	28%	27.6%	43.3%	***
	Large Adult Family	12%	12.3%	18.7%	***
	Urban	11%	11%	24%	
	Suburban	44%	44.5%	38.0%	*
	Rural	22%	22.0%	24.0%	
	Passive Smoking in household	22.9%	22.7%	31.3%	***

Notes: This table provides the summary statistics of the variables used in our. Column one displays the statistics for the overall sample, column two for households in which both parents are natural, column three for families with adoptive parents, and, finally, column four indicates the level of significance of the difference in means between households with natural parents and those with adoptive parents. The vertical panels shows first variables reflecting the characteristics of the child including overweight; second the parental overweight; third parental characteristics; and finally, other household characteristics. The level of significance of the t-test are indicated by the number of stars: * p<0.05 ** p<0.01*** p<0.001.

Table 2: Probit Model of the influence of parents being overweight on the likelihood of child being overweight

	(1)	(2)	(3)	(4)	(5)	(6)
Dependent variable:	Overweight		Overweight		Obese	
Control for parents being:	Overweight		Obese		Obese	
Type of Household	Both parents natural (13536)	Both parents adoptive (300)	Both parents natural (13536)	Both parents adoptive (300)	Both parents natural (13536)	Both parents adoptive (300)
Both	0.270*** (0.014)	0.210** (0.086)	0.342*** (0.019)	0.216* (0.129)	0.170*** (0.016)	0.208** (0.100)
Mum Only	0.129*** (0.019)	0.102 (0.124)	0.176*** (0.013)	-0.007 (0.068)	0.070*** (0.009)	- (0.009)
Dad Only	0.116*** (0.015)	0.240** (0.104)	0.144*** (0.013)	0.011 (0.068)	0.044*** (0.007)	0.025 (0.026)
Girl	0.047*** (0.007)	0.036 (0.047)	0.048*** (0.007)	0.046 (0.051)	0.016*** (0.004)	0.003 (0.010)
Observations	15534	15536	15534	15536	15534	15536
Log Likelihood	-6995.728	-157.207	-6995.551	-158.557	-2761.131	-48.932

Notes: This table reports the estimates of the probit models estimating the effect of measures of parental overweight on the likelihood of a child being overweight based on BMI. The rows identify the effect of both parents being overweight, only the mum being overweight or only the father being overweight. Given that gender might exert a specific effect, we include the effect of the child being a girl. The first column shows the effect of parental overweight on likelihood of the child being overweight when both parents are natural. The second column estimates the same for the sample of households when both parents are adoptive. In the third and fourth columns, we examine the effect for both household samples of parental obesity on child overweight. Finally, the last two columns estimate the effect of parental obesity on child obesity. Due to the reduced sample size, the last column does not produce estimates for the mother being obese. All estimates are marginal effects. The models control also for ethnicity, parents' education, passive smoking, flat ownership, and income. We provide robust standard errors in brackets.

Table 3: Models of the influence of parents overweight on child overweight correcting from sample selection bias of type of household

	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
Dependent variable:	Overweight			Overweight			Obese		
Control for parents being:	Overweight			Obese			Obese		
Type of Household	Both parents natural (13536)	Both parents adoptive (300)		Both parents natural (13536)	Both parents adoptive (300)		Both parents natural (13536)	Both parents adoptive (300)	
Model:	Heckprobit		PSM - correction	Heckprobit		PSM - correction	Heckprobit		PSM - correction
Both	0.252*** (0.015)	0.246** (0.086)	0.287*** (0.031)	0.277*** (0.016)	0.148** (0.119)	0.377*** (0.042)	0.102*** (0.011)	0.027*** (0.123)	0.181*** (0.038)
PSM interaction with Both			-0.781 (1.244)			-1.354 (1.360)			-0.171 (0.547)
Mother	0.115*** (0.016)	0.157 (0.129)	0.154*** (0.044)	0.154*** (0.011)	-0.014 (0.063)	0.166*** (0.031)	0.058*** (0.008)	-0.188 (0.811)	0.082*** (0.022)
PSM interaction with Only Mother			-0.935 (1.526)			0.222 (1.043)			-0.359 (0.481)
Father	0.109*** (0.014)	0.272*** (0.100)	0.106*** (0.033)	0.129*** (0.011)	-0.001 (0.061)	0.139*** (0.029)	0.039*** (0.007)	0.008 (0.039)	0.051*** (0.018)
PSM interaction with Only Father			0.507 (1.300)			0.077 (1.021)			-0.221 (0.495)
Girl	0.046*** (0.007)	0.046 (0.057)	0.046*** (0.007)	0.047*** (0.007)	0.038 (0.055)	0.048*** (0.007)	0.017*** (0.004)	0.002 (0.012)	0.015*** (0.004)
Observations	15534	15536	13826	15534	15536	13826	15534	15536	13826
Log Likelihood	-1.27e+04	-1574.124	-7166.998	-1.27e+04	-1575.787	-7170.681	-8451.805	-1456.645	-2823.569

Notes: In this table we report the estimates of the effect of parental overweight on the likelihood of a child being overweight (based on BMI) controlling for sample selection bias using two approaches, a Heckprobit and a Propensity Score Matching (PSM) specifications. As in Table 2, rows identify the both parents being overweight, only the mum being overweight or only the father being overweight. The coefficients in the additional row below each of these regressors reproduce the estimates of the interaction of the PSM indicator variable with both parents, only the mother or only the father being overweight. Again, we include the effect of the child being a girl. The first column shows the effect of parental overweight on likelihood of the child being overweight when both parents are natural. The second and third columns show the estimates for the sample of households when both parents are adoptive: Column two contains the marginal effects of the Heckprobit model and column three those correcting using the PSM specification. The fourth, fifth and sixth columns present the effect for both household samples of parental obesity on child overweight. The last three columns estimate the effect of parental obesity on child obesity. All estimates are marginal effects. The models control also for ethnicity, parents' education, passive smoking, flat ownership, and income. In the Heckprobit selection equation, we include parents' age, the father being unemployed or working full-time, mother's qualifications, type of household, and living in an urban area. Propensity score to be adopted based on gender of the child, mother working full-time, father being unemployed, household size, urban setting. We provide robust standard errors in brackets.

Table 4: Probit Models controlling for mother working full time

	(1)	(2)	(3)	(4)	(5)	(6)
Dependent variables:	Overweight		Overweight		Obese	
Control for parents being:	Overweight		Obese		Obese	
Type of Household	Both parents natural (13528)	Both parents adoptive (300)	Both parents natural (13528)	Both parents adoptive (300)	Both parents natural (13528)	Both parents adoptive (300)
Both	0.273*** (0.017)	0.212*** (0.064)	0.325*** (0.032)	0.240* (0.140)	0.150*** (0.025)	0.124 (0.101)
Both*(mother work FT=1)	-0.005 (0.013)	-0.003 (0.073)	0.019 (0.031)	-0.030 (0.128)	0.012 (0.013)	0.036 (0.061)
Mother	0.116*** (0.026)	0.047 (0.154)	0.194*** (0.021)	-0.087 (0.129)	0.065*** (0.013)	
Mother (mother work FT=1)	0.017 (0.024)	0.082 (0.177)	-0.022 (0.019)	0.121 (0.211)	0.004 (0.010)	
Dad	0.124*** (0.019)	0.241** (0.107)	0.106*** (0.022)	-0.029 (0.084)	0.026** (0.011)	0.009 (0.029)
Dad*(mother work FT=1)	-0.011 (0.016)	-0.001 (0.087)	0.047** (0.023)	0.071 (0.136)	0.020* (0.012)	0.020 (0.049)
Female	0.047*** (0.007)	0.035 (0.045)	0.048*** (0.007)	0.047 (0.047)	0.016*** (0.004)	0.004 (0.010)
Observations	13828	300	13828	300	13828	238
Log Likelihood	-6995.026	-157.080	-6991.820	-158.157	-2758.701	-48.441

Notes: This table reports the estimates of the probit models estimating the effect of measures of parental overweight on the likelihood of a child being overweight (based on BMI) examining if the mother working full time compounds the effect of parental overweight. The rows identify the effect of both parents being overweight, only the mum being overweight or only the father being overweight. The extra rows below each of these indicators include interactions with the mother working full time. As in Table 2, the first column shows the effect of parental overweight on likelihood of the child being overweight when both parents are natural. The second column estimates the same for the sample of households when both parents are adoptive. In the third and fourth columns, we examine the effect for both household samples of parental obesity on child overweight. Finally, the last two columns estimate the effect of parental obesity on child obesity. Due to the reduced sample size, the last column does not produce estimates for the mother being obese. All estimates are marginal effects. The models control also for ethnicity, parents' education, passive smoking, flat ownership, and income. We provide robust standard errors in brackets.

Appendices:

Appendix A1: Alternative measures of parental obesity

For the parents, we also construct measures of obesity (or of increased health risks due to being overweight) based on the Waist to Hip (WHIP) ratio, and on the waist circumference. For WHIP, we use the classification suggested by the WHO report on obesity and risk of diabetes (1999) by which men are considered obese if their WHIP exceeds 0.95, and women are obese if it exceeds 0.8. With respect to the waist circumference, we follow the National Institute for Health and Clinical Excellence (NICE) Guidelines by which the risk of health problems for men are increased if their waist is above or equal to 94cm and for women if above 80cm (Townsend et al, 2009). These two additional measures of obesity/increased risk of health problems due to excessive weight are used in our robustness checks. For children, the non-BMI-based alternative measures are not feasible as only 13.5% of all children in our sample have valid measures for waist and waist to hip ratios and an insignificant number of those not living with both their natural parents.

The simple probit models that use different measures of obesity show that the coefficient for both parents' being obese is not significant; that of the mother being obese is positive and significant for both measures (0.231 and 0.483, respectively); and that of the dad being obese is also significant and positive for the first measure WHIP. For the adopted children, the coefficients are not significant. We interpret this lack of results based on alternative measures as not surprising given that we are using different measures of obesity for children and their parents.

Appendix A2: Models with one biological parent - Mixed families

For mixed families, the effect of both parents being overweight or obese are positive and significant in all specifications, including the last three in which we control for the mother working full-time. The coefficient associated to only the mother being obese on the likelihood of the child being overweight is also always significant and positive. That only the dad is obese or overweight does not have a significant positive effect on the likelihood of the child being obese or overweight except

for the PSM-based estimates. The latter is possibly due to the fact that mixed families tend to have natural mothers and non-biological dads. When we control for the mother working full-time by interacting it with the overweight and obesity measures of the parents, the only remarkable effect is that it decreases slightly the transmission of obesity from both parents to children (column 12, coefficient -0.038).

Appendix A3: Correlation of parental and children's BMIs using OLS and Quantile Regression

The effect of the BMI of the mother on the BMI child is about 0.151 when both parents are natural and about 0.139 when they are mixed. The effect is not significant for the adopted group, possibly because of sample size issues. The effect of the father's BMI on the child's BMI is again significant and positive for natural parents' families and mixed (0.161 for the first group and 0.082 for the mixed one).

The quantile regression estimates for the 75% percentile for the BMI shows that for the upper tail of the BMI distribution, these effects are only strengthened, the effects of the mother's BMI are 0.213 and 0.180 for the both natural parents' and mixed families, respectively. The effect of the father's BMI is 0.223 and 0.082 for natural and mixed families, respectively. Being a female has a very large and significant impact on these two types of families too, being the coefficient 0.0389 and 0.569 for the general OLS specification but jumping to 0.611 and 0.807 respectively for the BMI upper 75% percentile.

