



Landscape risk analysis of pesticides applied in rice paddy areas in Europe: a case study for a watershed in Lombardy

Dimitrios G. Karpouzas,¹ C. Riparbelli,² E. Capri³

¹University of Thessaly, Greece

²ERSAF, Italy

³Universita Cattolica del Sacro Cuore, Italy



Marie Curie Fellowships

dkarpouz@agro.auth.gr


Πανεπιστήμιο Θεσσαλίας



Risk Assessment in rice in EU

“ Council and Commission recognized that due to particular conditions associated to rice cultivation and the specific criteria and principles referred to in Annex IV were inappropriate”

Tiered Risk Assessment Scheme in Rice




Mathematical Models for higher tier exposure assessment in rice (Tiers 3 and 4)

- **RICEWQ 1.6.4v:** PECs in paddy water, paddy sediment and PECs GW when used in an interface with the VADOFT sub-model (Tier 3)
- **RIVWQ 2.02v:** Surface water model used in combination with RICEWQ for receiving water systems (Tier 4)


Tier 4 - Basin Scale Exposure Assessment

Rice paddies, drainage and irrigation canals **and** related natural surface water bodies like streams and rivers **create a unique ecosystem which should be studied as a whole**





Spatial scaling up is relevant

Rice - area
Lombardy



Case study: pesticide exposure assessment for a small rice-cultivated watershed in Lombardy, Italy




Community of Tromello

What data were available?

Spatial data at scale 1:50,000 for the rice cultivated area of Lombardy (Lomellina), **Community of Tromello**


- **Technical card**
- **Aerial photographs**
- **Landuse**
- **Soil data (soil texture, OM, pH, permeability etc)**
- **Network of surface water bodies**



Tromello

Fiume Doppio (Torrente)

Light soils with high permeability




Teropopio (Torrente)


Soil Texture

- Loam
- Loamy Sand
- Sandy
- Sandy Loam

Surface water bodies in Lombardy



Lombardy



Province of Pavia

Availability of monitoring data for herbicides propanil and molinate at points just before and after the simulated watershed

Aim of the work

- Use of available GIS spatial data in combination with RICEWQ and RIVWQ models to assess the exposure of groundwater and surface water systems (**river Terdoppio**) to pesticides (**propanil and molinate**) used in a rice cultivated watershed in the community of Tromello, Lombardy

Studied Rice Cultivated Watershed

Total Watershed Area: 467 ha (201 paddy fields)

Pesticide Treatment (Scouting and interviews with farmers)

➤ **Molinate treated area: 73 ha (31 fields)**

➤ **Propanil treated area: 243.4 ha (85 fields)**

➤ **Area untreated or treated with other herbicides: 150.6 ha**

Mode of Cultivation (Local experts and interviews)



➤ **Traditional: 326.9 ha**

➤ **Dry – seeded: 140.1 ha**

Mode of Irrigation – Drainage

➤ **Continuous Flow-through system**

Paddy Pesticide Treatments



Surface water systems simulated

20 Drainage Canals



6 Streams

C. Brielli, R. Molinara, R.
Arcibalda, C. Magnani, C.
Ceccone, C. Dassi




River Terdoppio



Surface water bodies network

-  river terdoppio
-  drainage canals
-  streams



PECs calculation

**RICEWQ 10-year simulation in
201 paddies of the simulated
watershed**



PECs GW for year 1999



**Overflow/Runoff/Controlled
Drainage outputs of
RICEWQ used as inputs for
RIVWQ**



**RIVWQ simulation of the rice-cultivated watershed with 20
drainage canals linked to 6 streams which discharge their
water into river Terdoppio**



**PECs SW on the simulated river
Terdoppio for the year 1999**


RICEWQ parameterization

- **Irrigation/drainage and crop practices:** Local experts
- **Pesticide Treatment:** Scouting and interviews with farmers
- **Mode of Application:** Local experts
 - No drift for **molinate (granular application)**
 - Drift losses 2.77% of the application rate for **propanil**
- **Frequency of application:** 1 per year for molinate (4.5 kg/ha)
2 per year for propanil (4 kg/ha)
- **Pesticide Properties (K_{deg} , K_d , S_w):** Literature studies in aerobic water/sediment systems
- **Soil – Hydrology (percolation rate, soil texture):** Spatial GIS data

RIVWQ parameterization

Characteristics	Drainage Canals	Streams	River Terdoppio
Shape	Rectangular	Rectangular	Rectangular
Length (km)	Variable	Variable	7
Top Width (m)	1.5	6	15
Depth (m)	1.5	2	4
Flow velocity (cm/d)	0.01	0.07	0.2


GW Risk Assessment - Molinate



Validation – Surface water predictions


- Comparison of **PECs for propanil and molinate** at the point 6.5 km along the length of the simulated part of the river Terdoppio for the year 1999 with **measured values** at approximately the same point of the river
- Measured data were kindly provided by the Analytical Laboratory of the Province of Pavia

Validation - molinate



Generally good agreement between simulated and measured concentrations of molinate also at temporal level

Validation - propanil



Generally good agreement between simulated and measured concentrations of propanil

Closing remarks

- Propanil entails low risk for GW contamination unlike molinate whose PECs exceeded 0.1 µg/L in certain fields
- Good agreement between measured and simulated values of molinate and propanil in the river Terdoppio
- Landscape risk assessment using GIS as pre- or post-processing tool in combination with RICEWQ, RIVWQ models could be used in higher tier exposure analysis in rice cultivated basins

More information in our website

www.ricewq.org



The screenshot shows the homepage of the RiceWQ.org website. At the top, there is a green header with the logo "RiceWQ.org" and the tagline "pesticide risk assessment in rice". To the right of the logo are the European Union flag and the Marie Curie Fellowship logo. Below the header is a navigation menu with links: Home / [Who are we](#) / [Results](#) / [Presentations](#) / [Publications](#) / [Link & docs](#) / [TrainRice](#) / [Images](#). The main content area features a large green image of rice plants on the left. The title "What is RiceWQ" is followed by a paragraph: "Development and validation of tools for the establishment of a uniform European approach, in accordance to FOCUS scenarios, for [pesticide risk assessment in rice paddies](#)." Below this is the text "This goal will be achieved through..." followed by a numbered list of three items: 1. the development of well-defined national scenarios for rice cultivation at field scale; 2. the validation of appropriate mathematical models like [RICEWQ 1.6.2v](#) for the higher tier risk assessment in rice paddies in Europe; 3. the development of appropriate schemes for calculating the pesticide exposure resulted from rice paddies at the watershed-scale level; At the bottom of the page, there is a green footer with the text "RiceWQ - pesticide risk assessment in rice" and a small graphic of rice plants.

RiceWQ.org
pesticide risk assessment in rice

Home / [Who are we](#) / [Results](#) / [Presentations](#) / [Publications](#) / [Link & docs](#) / [TrainRice](#) / [Images](#)

What is RiceWQ

Development and validation of tools for the establishment of a uniform European approach, in accordance to FOCUS scenarios, for [pesticide risk assessment in rice paddies](#).

This goal will be achieved through...

- 1 the development of well-defined national scenarios for rice cultivation at field scale;
- 2 the validation of appropriate mathematical models like [RICEWQ 1.6.2v](#) for the higher tier risk assessment in rice paddies in Europe;
- 3 the development of appropriate schemes for calculating the pesticide exposure resulted from rice paddies at the watershed-scale level;

RiceWQ - pesticide risk assessment in rice

*Special thanks to [Marco Pastori](#), *ERSAF* for providing assistance and GIS data and to [Ms Broglia](#), *Province of Pavia* for providing the monitoring data*