

# Spatial and temporal patterns of pesticide concentrations in streamflow, drainage and runoff in a small Swedish agricultural catchment

Maria Sandin, Nicholas Jarvis, Kristin Piikki, Kevin Bishop, Mats Larsbo & Jenny Kreuger

Swedish University of Agricultural Sciences (SLU)

*York, England 30 Aug - 1 Sep, 2017*



# Background


Diffuse losses of pesticides to surface waters often originate from small fractions of the agricultural landscape susceptible to fast flow processes, i.e. surface runoff and/or macropore flow to drains

Cost-effective mitigation requires identification of the main contributing areas and understanding of the relative importance of different transport pathways


Very few studies carried out under Swedish agro-environmental conditions

Monitoring study in a small Swedish agricultural catchment:

- > **Can spatial variation in pesticide concentrations in streamflow be related to variations in soil texture?**
- > **Relative importance of surface and subsurface transport pathways?**



# A small Swedish agricultural catchment with large variation in soil types



# Sampling of streamflow, drainage and surface runoff

Stream



Drains



Surface runoff



- Sampling May and June 2013 – 2015
- LC-MS/MS analyses of 99 compounds
- Pesticide use data from annual farmer interviews

# Main results: Pesticide occurrence in stream water

clay soils > mixed soil types >> sandy soils


## *Pesticides detected in stream water*

	Clay soils	Mixed soil types	Sandy soils
# compounds	30-35	9-24	1-2
Maximum concentration $\mu\text{g l}^{-1}$	0.57-55	0.10-0.88	0.002-0.003
Sum of concentrations	3.31-65.21	0.17-1.90	0.003-0.005
#compounds >WQO*	4-7	1-2	0


\*Swedish national Water Quality Objectives are used to evaluate the environmental quality of surface waters

# Last application:

2009 - 2 years before sampling    year before sampling    year of sampling




average treated area %




Pesticides with GUS-index above 1.5 were more frequently detected in stream flow (i.e. pesticides in the upper left of each diagram)

Also, pesticides applied on larger areas were more frequently detected. However, especially in the clay soil area also pesticides with lower GUS-index were detected (applies also to previously applied pesticides).

Last application before sampling on 14<sup>th</sup> May 2014





Pesticide concentrations larger in surface runoff than in drainage



Topography implies that surface runoff water did not reach the stream, but infiltrated locally (i.e. lack of surface connectivity)



Macropore flow to drains likely dominant transport pathway



# Summary and conclusions

Clay soils>Mixed soil types>>Sandy soils

Soil texture maps potentially useful tool

Transport times often long – temporary storage along pathways

Macropore flow to drains likely dominant transport pathway



# Thank you for listening!



We thank the Swedish farmers' foundation for agricultural research and Centre for Chemical Pesticides for financial support

Read more in:

Sandin et al. (2018) Spatial and temporal patterns of pesticide concentrations in streamflow, drainage and runoff in a small Swedish agricultural catchment. *Science of the Total Environment* 610-611, 623-634

Maria's thesis: Surface and subsurface transport pathways of pesticides to surface waters

<https://pub.epsilon.slu.se/14474/>

