

Spectrally accurate global-local gyrokinetic simulations of turbulence in tokamak plasmas

Authors: Denis A. St-Onge, Felix I. Parra, Michael Barnes

Affiliation: University of Oxford

The suppression of turbulence in fusion plasmas, crucial to the success of next-generation tokamaks such as ITER, depends on a variety of physical mechanisms including the shearing of turbulent eddies via zonal flow and possibly the generation of intrinsic rotation. The turbulence exhibits interesting features such as avalanche structures and self-organisation, and its absence is associated with the formation of internal transport barriers. In order to successfully capture all of these effects in gyrokinetic simulation, one may need to allow for the inclusion of global effects (such as radial profile variation), as well as other often-neglected effects that are small in ρ_* . A careful numerical treatment is necessary to ensure that both the global and local physics are calculated accurately at reasonable expense.

To that end, we develop a novel approach to gyrokinetics where multiple flux-tube simulations are coupled together in a way that consistently incorporates global profile variation while allowing the use of Fourier basis functions, thus retaining spectral accuracy. By doing so, the need for Dirichlet boundary conditions typically employed in global gyrokinetic simulation, where fluctuations are nullified at the simulation boundaries, is obviated. This results in a smooth convergence to the local periodic limit as $\rho_* \rightarrow 0$. In addition, our scale-separated approach allows the use of transport-averaged sources and sinks, offering a more physically motivated alternative to the standard sources based on Krook-type operators.

Having implemented this approach in the flux-tube code `stella` [2], we study the role of transport barriers and avalanche formation in the transition region between the quiescent core and the turbulent pedestal, as well as the efficacy of intrinsic momentum generation by radial profile variation. Finally, we show that near-marginal plasmas can exhibit a radially localized Dimits shift [3], where strong coherent zonal flows give way to flows which are more turbulent and smaller scale.

This work has been carried out within the framework of the EUROfusion Consortium and has received funding from the Euratom research and training programme 2014–2018 and 2019–2020 under Grant Agreement No. 633053, and from the RCUK Energy Programme [Grant Number EP/P012450/1]. The views and opinions expressed herein do not necessarily reflect those of the European Commission. The authors acknowledge EUROfusion and the EUROfusion High Performance Computer (Marconi-Fusion).

References

- [1] F.I. Parra, M. Barnes. *Plasma Phys. Controlled Fusion*, 57(5):054003, 2015.
- [2] M. Barnes, F.I. Parra, M. Landreman. *J. Plasma Phys.*, 391:365, 2019.
- [3] A. M. Dimits, G. Bateman, *et al.* *Phys. Plasmas*, 7(3):969, 2000.